

## Self-Adapted Floquet Dynamics of Ultracold Bosons in a Cavity

Xi-Wang Luo and Chuanwei Zhang\*

*Department of Physics, The University of Texas at Dallas, Richardson, Texas 75080-3021, USA*

 (Received 19 December 2017; revised manuscript received 4 May 2018; published 29 June 2018)

Floquet dynamics of a quantum system subject to periodic modulations of system parameters provides a powerful tool for engineering new quantum matter with exotic properties. While system dynamics is significantly altered, the periodic modulation itself is usually induced externally and independent of Floquet dynamics. Here we propose a new type of Floquet physics for a Bose-Einstein condensate (BEC) subject to a shaken lattice generated inside a cavity, where the shaken lattice and atomic Floquet bands are mutually dependent, resulting in self-adapted Floquet dynamics. In particular, the shaken lattice induces Floquet quasienergy bands for the BEC, whose backaction leads to a self-adapted dynamical normal-superradiant phase transition for the shaken lattice. Such self-adapted Floquet dynamics shows two surprising and unique features: (i) The normal-superradiant phase transition possesses a hysteresis even without atom interactions. (ii) The dynamical atom-cavity steady state could exist at free energy maxima. The atom interactions strongly affect the phase transition of the BEC from zero to finite momenta. Our results provide a powerful platform for exploring self-adapted Floquet physics, which may open an avenue for engineering novel quantum materials.

DOI: [10.1103/PhysRevLett.120.263202](https://doi.org/10.1103/PhysRevLett.120.263202)

*Introduction.*—Floquet physics has been extensively studied in solid-state, ultracold atomic, and photonic systems in recent years with significant theoretical and experimental progress [1–14]. In particular, ultracold atoms in periodically driven optical lattices provide a highly controllable and disorder-free platform for studying Floquet physics, yielding many interesting and important phenomena such as coherent ac-induced tunneling and band coupling [15–25], the realization of gauge fields and topological bands [26–37], and the dynamical control of the expansion and quantum phase transition of bosonic systems [38–41], etc. These previous studies on Floquet physics assumed that system parameter modulations (e.g., the shaking or moving optical lattices) are determined solely by external driving and do not depend on system dynamics, i.e., no backaction of system Floquet states on parameter modulations. Therefore, a natural and important question is whether novel Floquet physics can emerge when system Floquet dynamics and parameter modulations are mutually dependent.

In this Letter, we address this important question by studying Floquet dynamics of ultracold boson atoms subject to a shaken optical lattice generated inside an optical cavity. In the past several decades, the interactions between atoms and static cavity fields with atom backactions (no Floquet physics) have been well studied in both theory [42] and experiment [43–53], showcasing rich cavity quantum electrodynamics (QED) physics ranging from few-body problems such as the Jaynes-Cummings model [54] to many-body physics such as the Dicke superradiance [55,56]. However, in these studies, the cavity

mode is static without periodic modulations such as shaking or moving.

Here we propose to realize a cavity-mode-induced, freely evolving shaken lattice, utilizing transverse pumping and a periodic modulation of the cavity field phase, and study its mutual interaction with a non- or weakly interacting Bose-Einstein condensate (BEC) inside the cavity. While such a shaken lattice generates Floquet bands for the BEC, the backaction of atom Floquet bands modulates the shaken lattice, leading to a dynamical superradiant phase, where atom Floquet bands and shaken lattice are self-adapted. Such Floquet normal-superradiant phase transition can be dramatically different from the non-Floquet one because of the coupling between different Floquet sidebands. In particular, the interplay between intra- and intersideband couplings may induce a hysteresis for the Floquet normal-superradiant phase transition of noninteracting atoms, yielding a completely new mechanism different from the well-known interaction-driven hysteresis [57–62]. Surprisingly, the steady state of the atom-cavity system can stabilize at the free energy maximum for dominant intersideband coupling because of the nonequilibrium nature of Floquet states. With increasing superradiant field, the Floquet band dispersion gradually evolves from a single minimum to doubly degenerate minima, leading to a second-order phase transition of the BEC. Such a transition can be significantly affected by the interaction between atoms through the backaction, which changes the critical superradiant field, the Floquet band dispersion, and the condensate momenta across the transition.




FIG. 1. (a) Schematic of the experimental setup. The cavity is pumped by an external transverse laser (red block arrow), with the cavity mode (red line) shaken by the two EOMs or PZT-driving mirrors. A BEC is prepared inside the cavity with a background lattice (blue line) generated by additional lasers (blue block arrows). The shaken cavity mode may induce interband couplings (dotted arrows) when the band gap matches the shaking frequency. (b) Energy levels of the atom and detunings of the cavity mode and pumping laser.

*System.*—As shown in Fig. 1(a), two schemes (A and B) can be used to generate a shaking cavity mode as  $\hat{\mathbf{z}}E(x, \varphi) \propto \hat{\mathbf{z}} \cos[k_0x + \varphi(t)]$ , with  $\hat{\mathbf{z}}$  being the polarization direction and  $k_0$  the wave number. Scheme A employs two electro-optic modulators (EOMs) [63], while scheme B uses two mirrors synchronously driven by piezoelectric transducers (PZTs) [64] to periodically and slowly [compared with the cavity free spectral range (FSR)] modify the optical path length does not change; therefore, the cavity resonance frequency is not affected. We consider a quasi-one-dimensional (the dynamics in other directions are reduced by a deep harmonic trap) BEC prepared in such a cavity, which is pumped by an external transverse laser. The pumping frequency  $\omega_p$  is close to the cavity resonance frequency  $\omega_c$ , both of which are detuned far below the atomic transition frequency  $\omega_a$  [see Fig. 1(b)].

After adiabatically eliminating the excited atomic level, we obtain the Hamiltonian of the atom-cavity system in a rotating frame with  $\hbar = 1$ :

$$\mathcal{H} = (\Delta_c - u)c^\dagger c + \int dx \Psi^\dagger(x) H_a(t) \Psi(x), \quad (1)$$

where  $c$  is the annihilation operator of the cavity photon, and  $\Psi(x)$  is the matter wave field of atoms in the ground state.  $\Delta_c = \omega_c - \omega_p$  is the cavity mode detuning, and  $u = (g_0^2/\Delta_e) \int dx \Psi^\dagger \Psi \cos^2(k_0x + \varphi)$  is the detuning induced by the atoms, which is typically small and negligible. The single-atom Hamiltonian is  $H_a(t) = -(\partial_x^2/2m) + V_{\text{ext}}(x) + \hat{V}_c(x, t)$ , with  $V_{\text{ext}}(x) = v_e \cos^2(k_0x)$  as a static external background lattice potential, which can be realized by additional lasers [65]. It gives rise to a static tight-binding atomic band structure  $\varepsilon_\lambda(q_x) = E_\lambda + t_\lambda \cos(q_x)$  with band index  $\lambda$  and Bloch momentum  $q_x \in [-k_0, k_0]$  [Fig. 2(a)].  $\hat{V}_c(x, t) = -\eta(c^\dagger + c/\sqrt{N_a}) \cos[k_0x + \varphi(t)]$  is the shaking potential induced by the cavity-assisted ac-Stark shift, with




FIG. 2. (a) Illustration of atomic static bands (solid lines) and phonon-dressed sidebands (dashed lines). (b) Couplings induced by  $\hat{V}_c$  in Eq. (2). The area of the Brillouin zone reduces in half by  $\hat{V}_c$ . The time-independent (time-dependent) term induces intrasideband (intersideband) couplings, indicated by solid arrows (dashed arrows).

$\eta = \Omega g_0 \sqrt{N_a} / \Delta_e$  being the coupling strength;  $\Delta_e = \omega_a - \omega_p$  the single-photon detuning;  $\Omega$  and  $g_0$  the Rabi frequencies of the transverse pumping laser and the single cavity photon, respectively ( $\Omega, g_0 \ll \Delta_e$ ); and the total atom number  $N_a$ .

Utilizing the expansion  $e^{if \cos \omega_0 t} = \sum_n i^n J_n(f) e^{in\omega_0 t}$  for  $\cos[k_0x + \varphi_0 + f \cos(\omega_0 t)]$ , we see that  $\hat{V}_c(x, t)$  can change the band structure by inducing a sequence of sidebands (i.e., phonon-dressed bands) and couplings between them (see Fig. 2). We choose  $v_e, \omega_0, \varphi_0$ , and  $f$  such that the static  $s$  band is near resonance with the two-phonon-dressed ( $\omega \equiv 2\omega_0$ )  $p$  band [Fig. 2(a)]; therefore, only these two bands need be considered for the calculation of the new Floquet band structure  $\tilde{\varepsilon}_\lambda(q_x)$ . The first-order expansion  $\propto e^{\pm i\omega_0 t}$  is far-off resonance and can be neglected. The zeroth-order expansion corresponds to intrasideband coupling; thus only terms containing  $\cos(k_0x)$  are nonzero due to the symmetry of the Wannier functions. Similarly, only terms with  $\sin(k_0x)$  are nonzero for the second-order expansion  $\propto e^{\pm i2\omega_0 t}$  that couples  $s$  and  $p$  bands. In total, the cavity-assisted potential can be written as [65]

$$\hat{V}_c = -\eta_0 \frac{c^\dagger + c}{\sqrt{N_a}} [\cos(k_0x) + 4\eta_1 \cos(\omega t) \sin(k_0x)], \quad (2)$$

where  $\eta_0 = \eta J_0(f) \cos(\varphi_0)$ , and  $\eta_1 = (J_2(f)/2J_0(f)) \tan(\varphi_0)$  (tunable by  $f$  and  $\varphi_0$ ) is the ratio between inter- and intrasideband coupling strengths. Note that the spatial period of  $\hat{V}_c(x, t)$  is twice that of  $V_{\text{ext}}(x)$ ; therefore the Brillouin zone (BZ) reduces by half to  $q_x \in [-k_0/2, k_0/2]$  through the band folding. Each band  $\lambda$  is split into two bands  $\lambda_0, \lambda_\pi$  [ $\lambda = s, p$ , as shown in Fig. 2(b)], and the lattice potential  $\hat{V}_c$  can only couple 0 and  $\pi$  bands due to the momentum transfer.  $\varphi_0$  characterizes the relative phase between background lattice and the shaking center of the cavity field: for  $\varphi_0 = 0$  ( $\pi/2$ ), the shaking potential is symmetric (asymmetric) at each site of the background lattice; therefore it can only induce intrasideband (intersideband) couplings. Both couplings coexist for  $\varphi_0 \neq j\pi/2$  ( $j$  is an integer) [65].

*Method.*—Under mean-field approximation, the cavity field satisfies the Langevin equation  $i\langle\dot{c}(t)\rangle = \langle[c, \mathcal{H}]\rangle - i\gamma/2\langle c(t)\rangle$  due to the weak leakage of the high- $Q$  cavity, yielding

$$i\dot{\alpha} = (\Delta_c - u - i\gamma/2)\alpha - \eta_0\Theta(t), \quad (3)$$

with  $\alpha(t) = \langle c(t)\rangle/\sqrt{N_a}$  and  $\gamma$  as the cavity loss rate. Here  $\Theta(t) = N_a^{-1} \int dx \langle \Psi^\dagger \Psi \rangle [\cos(k_0 x) + 4\eta_t \sin(k_0 x) \cos(\omega t)]$  is the atomic density order. The frequency  $\omega$  is chosen to be much larger than  $\eta_0\Theta$ ,  $\Delta_c - u$ ,  $\gamma/2$  (see the *Experimental consideration* section below), so that high-order oscillation terms  $\propto e^{\pm i\omega t}$  can be dropped in Eq. (3). The steady-state solution in the presence of cavity loss is determined by  $\dot{\alpha} = 0$ , yielding

$$\phi = 2\tilde{\Delta}_c\eta_0\Theta_0/(\tilde{\Delta}_c^2 + \gamma^2/4), \quad (4)$$

where  $\phi \equiv \alpha + \alpha^*$ ,  $\tilde{\Delta}_c = \Delta_c - u_0$  is the effective detuning with  $u_0 = (1/T) \int_0^T dt u(t)$  a small constant, and  $\Theta_0 = (1/T) \int_0^T dt \Theta(t)$  with  $T = 2\pi/\omega$  the period. The shaking cavity field  $\phi$  and atom density order  $\Theta_0$  are determined self-consistently through Eq. (4).

In the self-consistent determination, we replace  $(c^\dagger + c/\sqrt{N_a})$  in  $\hat{V}_c(x, t)$  [Eq. (2)] with  $\phi$  for the Floquet Hamiltonian  $H_a(t)$  [Eq. (1)] and find Floquet quasienergy bands and Floquet states for the BEC, from which the atom density order  $\Theta$  can be calculated [65].  $\Theta$  in turn drives the cavity field  $\phi$  through Eq. (3) as the atom feedback, yielding the self-adapted steady solution in Eq. (4). We find that solving the self-consistent equation in Eq. (4) is equivalent to finding the extremum of the free energy density  $F(\phi) = \langle \mathcal{H} \rangle / N_a$  (i.e.,  $(\partial F / \partial \phi) = 0$ ) [65]. Notice that  $\phi = \Theta_0 = 0$  is always a trivial solution. Across the transition from normal to superradiant phases, the zero solution becomes unstable and  $\phi$ ,  $\Theta_0$  evolve from zero to finite values.

*Results.*—We focus on noninteracting BECs and discuss the interaction effects later. The numerically calculated phase diagrams and corresponding self-adapted Floquet bands with  $\omega \gtrsim E_{\text{sp}} + |t_s| + |t_p|$  are shown in Fig. 3. For a small  $\eta_t$  ( $\lesssim 1$ ),  $\hat{V}_c$  is dominated by the time-independent term that couples  $s_0$  and  $s_\pi$  bands, which would lower the energy of the  $s_0$  band [see the inset in Fig. 3(a)], indicating  $\langle \hat{V}_c \rangle = -\eta_0\phi\Theta_0 < 0$ . According to Eq. (4), a nontrivial steady-state solution exists only for blue effective detuning  $\tilde{\Delta}_c > 0$  [Figs. 3(a) and 3(b)]. The phase transition requires a stronger  $\eta_0$  as  $\eta_t$  increases, indicating that the transition becomes harder due to the competition between inter- and intrasideband couplings. As  $\tilde{\Delta}_c$  decreases, the critical value of  $\eta_0$  required for superradiance first decreases, then increases and tends to infinity at  $|\tilde{\Delta}_c| = 0$ . This is because  $\hat{V}_c$ , which drives the atomic density order  $\Theta_0$ , is proportional to  $\phi$  and approaches zero as  $|\tilde{\Delta}_c| \rightarrow 0$  [see Eq. (4)].




FIG. 3. (a), (b) Phase diagrams for dominant intrasideband coupling, with (a)  $\tilde{\Delta}_c = 1$  and (b)  $\eta_t = 0.5$ . I (II) represents the normal (superradiant) phase. (c), (d) Phase diagrams for dominant intersideband coupling, with (c)  $\tilde{\Delta}_c = -1$  and (d)  $\eta_t^{-1} = 0.1$ ,  $\eta_{0t} = \eta_0\eta_t$ . III represents the hysteresis region. The insets in (a) and (c) show the  $s_0$ -band dispersion in the normal (red dashed line) and superradiant phases (blue solid line). The three blue (red) dots in (b) [(d)] belong to different phases, and they mark the points where the free energy curves are plotted in the inset. The parameters are  $v_e = -6$ ,  $\omega = 4.8$ , and  $\gamma = 2$  (with energy unit  $E_R = \hbar^2 k_0^2 / 2m$ ), and atom-atom interaction is neglected.

We find that the solution is located at the minima of  $F(\phi)$ , which can be expanded as  $F(\phi) = a_2\phi^2 + a_4\phi^4 + \dots$ .  $F(\phi)$  exhibits a continuous transition from a single minimum at  $\phi = 0$  to double minima at  $\phi \neq 0$  [see the inset in Fig. 3(b)], where  $a_2$  and  $a_4$  are both positive before the transition, and  $a_2$  changes sign when the phase transition (second order) occurs.

For a large  $\eta_t$  ( $\gg 1$ ),  $\hat{V}_c$  is dominated by the interband coupling between the  $s_0$  band and  $p_\pi$  band that has a lower energy than the  $s_0$  band; therefore, atoms stay at the high-energy excited band, and increasing the cavity field would raise the band energy [the inset in Fig. 3(c)] of the BEC, leading to  $\langle \hat{V}_c \rangle = -\eta_0\phi\Theta_0 > 0$ . As a result, the nontrivial steady-state solution exists only for red effective detuning  $\tilde{\Delta}_c < 0$  [Figs. 3(c) and 3(d)]. Surprisingly, the steady-state solution is found at the maxima of  $F(\phi)$ , which exhibit a transition from a single maximum at  $\phi = 0$  to double maxima at  $\phi \neq 0$  [the inset in Fig. 3(d)] because of the nonequilibrium phase transition of the dynamical steady states which may not minimize the energy. Without Floquet sidebands, such superradiance at energy maximum for  $\tilde{\Delta}_c < 0$  would not exist, because atoms generally prefer staying in the lowest static band which can only couple with higher static bands.

Moreover, depending on the value of  $\eta_t$ , the transition can be either continuous (second order) or discontinuous with a hysteresis loop (first order) [Fig. 4(b)]. Such hysteresis originates from the interplay between intrasideband and intersideband couplings, which induces a second-order coupling (similar to a two-photon Raman process)




FIG. 4. (a), (c) Superradiant order parameters and condensate momenta versus  $\eta_0$  and  $\eta_0 t$ . Stars and circles mark the transition points between zero and finite condensate momenta for  $g = 0$  and  $g > 0$ , respectively. The inset of (a) shows the band structure for  $|q_x^m| > 0$ . A clear hysteresis loop appears in (c) with solid lines (dashed lines) corresponding to stable (unstable) steady-state solutions. (b), (d) Difference between  $q_x^c$  (red dotted line) and  $q_x^m$  (blue dashed line) for  $g > 0$ . In the gray region, bosons are condensed into the local band maximum at  $q_x = 0$ . The black solid line shows  $q_x^m = q_x^c$  for  $g = 0$ . (a), (b) [(c), (d)] correspond to dominant intrasideband (intersideband) couplings with  $\Delta_c = 1$ ,  $\eta_l = 0.5$  and  $g\bar{n} = 0.02$  ( $\Delta_c = -1$ ,  $\eta_l^{-1} = 0.15$  and  $g\bar{n} = 0.1$ ).  $\bar{n} = N_a/L$  is the average density, with  $L$  as the system size. Other parameters are the same as in Fig. 3.

between the  $s_0$  and  $p_0$  bands mediated by the  $s_\pi$  and  $p_\pi$  bands [see Fig. 2(b)]. Notice that the  $p_0$  band is just below the  $s_0$  band; therefore, this second-order coupling raises the  $s_0$  band by increasing  $a_4$ . As a result,  $a_4$  may change sign (from negative to positive) prior to  $a_2$  changing sign (from negative to positive) when the second-order coupling is strong enough, leading to a multistability behavior where  $F(\phi)$  exhibits three maxima simultaneously.

Hysteresis phenomena are related to strong nonlinearities [78], which are usually induced by strong atom-atom interactions [57,60]. For example, the strong Ising interaction in the Dicke model can lead to a hysteresis loop of the superradiant phase transition [57–59]. However, the hysteresis effect in our system does not need atom-atom interaction at all, and has a completely different mechanism originating from the coupling between Floquet sidebands induced by the shaken cavity mode. Our study offers an excellent example and a realistic system for observing hysteresis effects without atom interactions.

As the superradiant field increases, the  $s_0 - p_0$  band coupling would induce a transition of the Floquet band  $\tilde{\epsilon}_{s_0}(q_x)$  from a single minimum at  $q_x = 0$  to doubly degenerate minima at  $q_x \neq 0$  [see inset in Fig. 4(a)]. In our system, the  $s_0 - p_0$  band coupling is a Raman-like coupling mediated by  $s_\pi$  and  $p_\pi$ ; therefore such a transition should be observed when the inter- and intrasideband couplings coexist. We consider atom-atom interaction (tunable through Feshbach resonance [79]) that is weak and repulsive; therefore the interaction energy is always minimized when the system stays in a single momentum

state. We assume that  $\tilde{\epsilon}_{s_0}(q_x)$  is minimized at  $\pm q_x^m$ . For negligibly weak interaction, the condensate momentum  $q_x^c$  would locate at one of the band minima (either  $+q_x^m$  or  $-q_x^m$ ) due to spontaneous symmetry breaking [80]. Such a transition in the BEC would also lead to a second-order transition [stars in Figs. 4(a) and 4(c)] of the shaking cavity field  $\phi_s$  due to the backaction.

As the atom-atom interaction increases [still weak enough such that the superradiant phase transition is not affected and the long-time behaviors (heating, atom loss, etc.) are not significant [22,25,81]], the phase transition of the BECs from  $q_x^c = 0$  to  $q_x^c \neq 0$  may be shifted because  $q_x^c \neq \pm q_x^m$ .  $q_x^c$  should be determined by minimizing  $\tilde{\epsilon}_{s_0}(q_x) + \tilde{\epsilon}_{\text{int}}(q_x)$ , with  $\tilde{\epsilon}_{\text{int}}(q_x)$  being the momentum-dependent interaction energy,

$$\tilde{\epsilon}_{\text{int}}(q_x) = \frac{1}{T} \int dt \int dx \frac{g}{2} |n_{s_0, q_x}(x, t)|^2. \quad (5)$$

Here  $g$  is the interaction constant, and  $n_{s_0, q_x}(x, t) = \langle \Psi^\dagger \Psi \rangle$  is the atom density. In our system, band mixing enhances spatial modulation of the density, and the interaction energy  $\tilde{\epsilon}_{\text{int}}(q_x)$  is minimized at  $q_x = 0$ , where the mixing is the smallest. As a result,  $q_x^c$  is smaller than  $q_x^m$ , and bosons may be condensed into the local maximum of the single-particle band at  $q_x = 0$  [see Figs. 4(b) and 4(d)]. The transition from  $q_x^c = 0$  to  $q_x^c \neq 0$  for  $g > 0$  requires a stronger superradiant field than the transition for  $g = 0$  [see Figs. 4(a) and 4(c)], and it also leads to a second-order transition of  $\phi_s$ , which in turn leads to a transition in  $q_x^m$ .

*Experimental consideration.*—We consider a high-finesse (low-loss) cavity with  $\gamma = 2E_R$  (with  $E_R = \hbar^2 k_0^2 / 2m$  as the recoil energy). Generally, atoms with a small mass are preferred to obtain a large  $E_R$ , thus a large  $\gamma$ , which makes the cavity easy to realize. For example,  $^7\text{Li}$  ( $^{23}\text{Na}$ ) atoms have a recoil energy  $E_R \sim 40$  kHz (10 kHz), corresponding to  $\gamma = 80$  kHz (20 kHz), which can be realized with current technique [65,82]. The shaking frequency  $\omega_0$  is about several tens of kilohertz for  $^{23}\text{Na}$  and several hundred kilohertz for  $^7\text{Li}$  (both much smaller than the free spectral range, which is in gigahertz), and such phase modulation can be implemented by PZT-driving mirrors or low-loss EOMs [65]. The system studied here only involves a change of introducing PZT-driving mirrors or inserting two EOMs into the setups used in the ETH and Hamburg laboratories [45,49,52,53,82], and thus should be feasible with current technology. Moreover, our model can also be implemented by combining a shaking external lattice and a periodic driving force (e.g., using a periodically modulated magnetic field gradient) [65].

*Discussion.*—We proposed a new type of Floquet physics, where the parameter modulations are not only related to external driving, but also mutually coupled with system dynamics. As an example, we studied such Floquet dynamics of BECs in shaking optical lattices, which lead

to interesting new phenomena including self-adapted shaking fields, Floquet bands, and hysteresis effects. Such self-adapted Floquet physics may also arise in various other systems such as Rydberg atom or molecule micromaser [83], ion-trap cavity [84], and circuit quantum electrodynamics (circuit QED) [85], etc. For example, in a circuit QED system, the electromagnetic field on the waveguide resonator can be periodically modulated by attaching superconducting quantum interference devices (SQUIDs) to the ends of the resonator, with the SQUIDs driving by external magnetic fluxes [86]. Such a modulated waveguide resonator can couple with superconducting qubits (artificial atoms), and these qubits may also strongly couple with each other, where interesting self-adapted Floquet dynamics may emerge. Such self-adapted Floquet circuit QED may find important applications in quantum information processing and will be addressed in future works. Within the shaking lattice cavity system, interesting physics may also arise by considering strong atom-atom interaction (Bose-Hubbard model [52,53]) or strong atom-single-photon coupling (limit cycles and chaos [87]), and the superradiance of fermion gases (topological bands [88,89]). In this spirit, our proposal opens up new possibilities for studying self-adapted Floquet physics in various systems, which may pave a way for engineering new exotic quantum matter.

We thank A. Hemmerich for helpful discussion. This work is supported by AFOSR (Grant No. FA9550-16-1-0387), the National Science Foundation (Grant No. PHY-1505496), and ARO (Grant No. W911NF-17-1-0128).

---

\*chuanwei.zhang@utdallas.edu

- [1] N. H. Lindner, G. Refael, and V. Galitski, Floquet topological insulator in semiconductor quantum wells, *Nat. Phys.* **7**, 490 (2011).
- [2] Y. H. Wang, H. Steinberg, P. Jarillo-Herrero, and N. Gedik, Observation of Floquet-Bloch states on the surface of a topological insulator, *Science* **342**, 453 (2013).
- [3] U. De Giovannini, H. Hübener, and A. Rubio, Monitoring electron-photon dressing in WSe<sub>2</sub>, *Nano Lett.* **16**, 7993 (2016).
- [4] J. Cayssol, B. Dóra, F. Simon, and R. Moessner, Floquet topological insulators, *Phys. Status Solidi RRL* **7**, 101 (2013).
- [5] H. Hübener, M. A. Sentef, U. De Giovannini, A. F. Kemper, and A. Rubio, Creating stable Floquet-Weyl semimetals by laser-driving of 3D Dirac materials, *Nat. Commun.* **8**, 13940 (2017).
- [6] N. Goldman and J. Dalibard, Periodically Driven Quantum Systems: Effective Hamiltonians and Engineered Gauge Fields, *Phys. Rev. X* **4**, 031027 (2014).
- [7] M. Holthaus, Floquet engineering with quasienergy bands of periodically driven optical lattices, *J. Phys. B* **49**, 013001 (2016).
- [8] A. Eckardt, Colloquium: Atomic quantum gases in periodically driven optical lattices, *Rev. Mod. Phys.* **89**, 011004 (2017).
- [9] T. Kitagawa, M. A. Broome, A. Fedrizzi, M. S. Rudner, E. Berg, I. Kassal, A. Aspuru-Guzik, E. Demler, and A. G. White, Observation of topologically protected bound states in photonic quantum walks, *Nat. Commun.* **3**, 882 (2012).
- [10] K. Fang, Z. Yu, and S. Fan, Realizing effective magnetic field for photons by controlling the phase of dynamic modulation, *Nat. Photonics* **6**, 782 (2012).
- [11] M. C. Rechtsman, J. M. Zeuner, Y. Plotnik, Y. Lumer, D. Podolsky, F. Dreisow, S. Nolte, M. Segev, and A. Szameit, Photonic floquet topological insulators, *Nature (London)* **496**, 196 (2013).
- [12] K. Fang, Z. Yu, and S. Fan, Photonic Aharonov-Bohm Effect Based on Dynamic Modulation, *Phys. Rev. Lett.* **108**, 153901 (2012).
- [13] E. Li, B. J. Eggleton, K. Fang, and S. Fan, Photonic Aharonov-Bohm effect in photon-phonon interactions, *Nat. Commun.* **5**, 3225 (2014).
- [14] D. N. Basov, R. D. Averitt, and D. Hsieh, Towards properties on demand in quantum materials, *Nat. Mater.* **16**, 1077 (2017).
- [15] D. H. Dunlap and V. M. Kenkre, Dynamic localization of a charged particle moving under the influence of an electric field, *Phys. Rev. B* **34**, 3625 (1986).
- [16] V. V. Ivanov, A. Alberti, M. Schioppo, G. Ferrari, M. Artoni, M. L. Chiofalo, and G. M. Tino, Coherent Delocalization of Atomic Wave Packets in Driven Lattice Potentials, *Phys. Rev. Lett.* **100**, 043602 (2008).
- [17] C. Sias, H. Lignier, Y. P. Singh, A. Zenesini, D. Ciampini, O. Morsch, and E. Arimondo, Observation of Photon-Assisted Tunneling in Optical Lattices, *Phys. Rev. Lett.* **100**, 040404 (2008).
- [18] E. Haller, R. Hart, M. J. Mark, J. G. Danzl, L. Reichsöllner, and H.-C. Nägerl, Inducing Transport in a Dissipation-Free Lattice with Super Bloch Oscillations, *Phys. Rev. Lett.* **104**, 200403 (2010).
- [19] A. Alberti, V. V. Ivanov, G. M. Tino, and G. Ferrari, Engineering the quantum transport of atomic wavefunctions over macroscopic distances, *Nat. Phys.* **5**, 547 (2009).
- [20] N. Gemelke, E. Sarajlic, Y. Bidel, S. Hong, and S. Chu, Parametric Amplification of Matter Waves in Periodically Translated Optical Lattices, *Phys. Rev. Lett.* **95**, 170404 (2005).
- [21] W. S. Bakr, P. M. Preiss, M. E. Tai, R. Ma, J. Simon, and M. Greiner, Orbital excitation blockade and algorithmic cooling in quantum gases, *Nature (London)* **480**, 500 (2011).
- [22] C. V. Parker, L.-C. Ha, and C. Chin, Direct observation of effective ferromagnetic domains of cold atoms in a shaken optical lattice, *Nat. Phys.* **9**, 769 (2013).
- [23] L.-C. Ha, L. W. Clark, C. V. Parker, B. M. Anderson, and C. Chin, Roton-Maxon Excitation Spectrum of Bose Condensates in a Shaken Optical Lattice, *Phys. Rev. Lett.* **114**, 055301 (2015).
- [24] Z. Zheng, C. Qu, X. Zou, and C. Zhang, Fulde-Ferrell Superfluids without Spin Imbalance in Driven Optical Lattices, *Phys. Rev. Lett.* **116**, 120403 (2016).
- [25] M. A. Khamehchi, C. Qu, M. E. Mossman, C. Zhang, and P. Engels, Spin-momentum coupled Bose-Einstein

- condensates with lattice band pseudospins, *Nat. Commun.* **7**, 10867 (2016).
- [26] A. S. Sørensen, E. Demler, and M. D. Lukin, Fractional Quantum Hall States of Atoms in Optical Lattices, *Phys. Rev. Lett.* **94**, 086803 (2005).
- [27] T. Kitagawa, E. Berg, M. Rudner, and E. Demler, Topological characterization of periodically driven quantum systems, *Phys. Rev. B* **82**, 235114 (2010).
- [28] M. Aidelsburger, M. Atala, M. Lohse, J. T. Barreiro, B. Paredes, and I. Bloch, Realization of the Hofstadter Hamiltonian with Ultracold Atoms in Optical Lattices, *Phys. Rev. Lett.* **111**, 185301 (2013).
- [29] M. Aidelsburger, M. Atala, S. Nascimbène, S. Trotzky, Y.-A. Chen, and I. Bloch, Experimental Realization of Strong Effective Magnetic Fields in an Optical Lattice, *Phys. Rev. Lett.* **107**, 255301 (2011).
- [30] J. Struck, C. Ölschläger, M. Weinberg, P. Hauke, J. Simonet, A. Eckardt, M. Lewenstein, K. Sengstock, and P. Windpassinger, Tunable Gauge Potential for Neutral and Spinless Particles in Driven Optical Lattices, *Phys. Rev. Lett.* **108**, 225304 (2012).
- [31] J. Struck, M. Weinberg, C. Ölschläger, P. Windpassinger, J. Simonet, K. Sengstock, R. Höppner, P. Hauke, A. Eckardt, M. Lewenstein, and L. Mathey, Engineering Ising-XY spin-models in a triangular lattice using tunable artificial gauge fields, *Nat. Phys.* **9**, 738 (2013).
- [32] H. Miyake, G. A. Siviloglou, C. J. Kennedy, W. C. Burton, and W. Ketterle, Realizing the Harper Hamiltonian with Laser-Assisted Tunneling in Optical Lattices, *Phys. Rev. Lett.* **111**, 185302 (2013).
- [33] M. Atala, M. Aidelsburger, M. Lohse, J. T. Barreiro, B. Paredes, and I. Bloch, Observation of chiral currents with ultracold atoms in bosonic ladders, *Nat. Phys.* **10**, 588 (2014).
- [34] C. J. Kennedy, W. C. Burton, W. C. Chung, and W. Ketterle, Observation of Bose-Einstein condensation in a strong synthetic magnetic field, *Nat. Phys.* **11**, 859 (2015).
- [35] M. E. Tai, A. Lukin, M. Rispoli, R. Schittko, T. Menke, D. Borgnia, P. M. Preiss, F. Grusdt, A. M. Kaufman, and M. Greiner, Microscopy of the interacting Harper-Hofstadter model in the two-body limit, *Nature (London)* **546**, 519 (2017).
- [36] G. Jotzu, M. Messer, R. Desbuquois, M. Lebrat, T. Uehlinger, D. Greif, and T. Esslinger, Experimental realization of the topological Haldane model with ultracold fermions, *Nature (London)* **515**, 237 (2014).
- [37] M. Aidelsburger, M. Lohse, C. Schweizer, M. Atala, J. T. Barreiro, S. Nascimbene, N. R. Cooper, I. Bloch, and N. Goldman, Measuring the Chern number of Hofstadter bands with ultracold bosonic atoms, *Nat. Phys.* **11**, 162 (2015).
- [38] A. Eckardt, C. Weiss, and M. Holthaus, Superfluid-Insulator Transition in a Periodically Driven Optical Lattice, *Phys. Rev. Lett.* **95**, 260404 (2005).
- [39] H. Lignier, C. Sias, D. Ciampini, Y. Singh, A. Zenesini, O. Morsch, and E. Arimondo, Dynamical Control of Matter-Wave Tunneling in Periodic Potentials, *Phys. Rev. Lett.* **99**, 220403 (2007).
- [40] A. Zenesini, H. Lignier, D. Ciampini, O. Morsch, and E. Arimondo, Coherent Control of Dressed Matter Waves, *Phys. Rev. Lett.* **102**, 100403 (2009).
- [41] J. Struck, C. Ölschläger, R. L. Targat, P. Soltan-Panahi, A. Eckardt, M. Lewenstein, P. Windpassinger, and K. Sengstock, Quantum simulation of frustrated classical magnetism in triangular optical lattices, *Science* **333**, 996 (2011).
- [42] H. Walther, B. T. H. Varcoe, B.-G. Englert, and T. Becker, Cavity quantum electrodynamics, *Rep. Prog. Phys.* **69**, 1325 (2006).
- [43] A. T. Black, H. W. Chan, and V. Vuletić, Observation of Collective Friction Forces due to Spatial Self-Organization of Atoms: From Rayleigh to Bragg Scattering, *Phys. Rev. Lett.* **91**, 203001 (2003).
- [44] S. Slama, S. Bux, G. Krenz, C. Zimmermann, and Ph. W. Courteille, Superradiant Rayleigh Scattering and Collective Atomic Recoil Lasing in a Ring Cavity, *Phys. Rev. Lett.* **98**, 053603 (2007).
- [45] K. Baumann, C. Guerlin, F. Brennecke, and T. Esslinger, Dicke quantum phase transition with a superfluid gas in an optical cavity, *Nature (London)* **464**, 1301 (2010).
- [46] K. Baumann, R. Mottl, F. Brennecke, and T. Esslinger, Exploring Symmetry Breaking at the Dicke Quantum Phase Transition, *Phys. Rev. Lett.* **107**, 140402 (2011).
- [47] K. J. Arnold, M. P. Baden, and M. D. Barrett, Self-Organization Threshold Scaling for Thermal Atoms Coupled to a Cavity, *Phys. Rev. Lett.* **109**, 153002 (2012).
- [48] H. Ritsch, P. Domokos, F. Brennecke, and T. Esslinger, Cold atoms in cavity-generated dynamical optical potentials, *Rev. Mod. Phys.* **85**, 553 (2013).
- [49] H. Keßler, J. Klinder, M. Wolke, and A. Hemmerich, Steering Matter Wave Superradiance with an Ultranarrow-Band Optical Cavity, *Phys. Rev. Lett.* **113**, 070404 (2014).
- [50] M. R. Bakhtiari, A. Hemmerich, H. Ritsch, and M. Thorwart, Nonequilibrium Phase Transition of Interacting Bosons in an Intracavity Optical Lattice, *Phys. Rev. Lett.* **114**, 123601 (2015).
- [51] S. Gopalakrishnan, B. L. Lev, and P. M. Goldbart, Emergent crystallinity and frustration with Bose-Einstein condensates in multimode cavities, *Nat. Phys.* **5**, 845 (2009).
- [52] J. Klinder, H. Keßler, M. Reza Bakhtiari, M. Thorwart, and A. Hemmerich, Observation of a Superradiant Mott Insulator in the Dicke-Hubbard Model, *Phys. Rev. Lett.* **115**, 230403 (2015).
- [53] R. Landig, L. Hruby, N. Dogra, M. Landini, R. Mottl, T. Donner, and T. Esslinger, Quantum phases from competing short-and long-range interactions in an optical lattice, *Nature (London)* **532**, 476 (2016).
- [54] B. W. Shore and P. L. Knight, The Jaynes-Cummings model, *J. Mod. Opt.* **40**, 1195 (1993).
- [55] R. H. Dicke, Coherence in spontaneous radiation processes, *Phys. Rev.* **93**, 99 (1954).
- [56] D. Nagy, G. Kónya, G. Szirmai, and P. Domokos, Dicke-Model Phase Transition in the Quantum Motion of a Bose-Einstein Condensate in an Optical Cavity, *Phys. Rev. Lett.* **104**, 130401 (2010).
- [57] C. F. Lee and N. F. Johnson, First-Order Superradiant Phase Transitions in a Multiqubit Cavity System, *Phys. Rev. Lett.* **93**, 083001 (2004).
- [58] S. Gammelmark and K. Mølmer, Phase transitions and Heisenberg limited metrology in an Ising chain interacting with a single-mode cavity field, *New J. Phys.* **13**, 053035 (2011).

- [59] X.-W. Luo, Y.-N. Zhang, X. Zhou, G.-C. Guo, and Z.-W. Zhou, Dynamic phase transitions of a driven Ising chain in a dissipative cavity, *Phys. Rev. A* **94**, 053809 (2016).
- [60] S. Eckel, J. G. Lee, F. Jendrzejewski, N. Murray, C. W. Clark, C. J. Lobb, W. D. Phillips, M. Edwards, and G. K. Campbell, Hysteresis in a quantized superfluid “atomtronic” circuit, *Nature (London)* **506**, 200 (2014).
- [61] C. Hamner, C. Qu, Y. Zhang, J. Chang, M. Gong, C. Zhang, and P. Engels, Dicke-type phase transition in a spin-orbit-coupled Bose-Einstein condensate, *Nat. Commun.* **5**, 4023 (2014).
- [62] A. Trenkwalder, G. Spagnolli, G. Semeghini, S. Coop, M. Landini, P. Castilho, L. Pezze, G. Modugno, M. Inguscio, A. Smerzi, and M. Fattori, Quantum phase transitions with parity-symmetry breaking and hysteresis, *Nat. Phys.* **12**, 826 (2016).
- [63] A. Yariv and P. Yeh, *Photonics: Optical Electronics in Modern Communications* (Oxford University Press, New York, 2007).
- [64] D. Goldovsky, V. Jouravsky, and A. Pe’er, Simple and robust phase-locking of optical cavities with  $> 200$  KHz servo-bandwidth using a piezo-actuated mirror mounted in soft materials, *Opt. Express* **24**, 28239 (2016).
- [65] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.120.263202> more details about the cavity-assisted potential, the Floquet bands and energy density, the stability analysis, and the experimental consideration, which includes Refs. [66–77].
- [66] R. Gehr, J. Volz, G. Dubois, T. Steinmetz, Y. Colombe, B. L. Lev, R. Long, J. Esteve, and J. Reichel, Cavity-Based Single Atom Preparation and High-Fidelity Hyperfine State Readout, *Phys. Rev. Lett.* **104**, 203602 (2010).
- [67] M. J. Bhaseen, J. Mayoh, B. D. Simons, and J. Keeling, Dynamics of nonequilibrium Dicke models, *Phys. Rev. A* **85**, 013817 (2012).
- [68] D. Nagy, G. Szirmai, and P. Domokos, Critical exponent of a quantum-noise-driven phase transition: The open-system Dicke model, *Phys. Rev. A* **84**, 043637 (2011).
- [69] T. Nakamura, S. Tani, I. Ito, and Y. Kobayashi, Beyond 500-kHz bandwidth piezo-electric transducers For GHz-Comb applications, *2017 European Conference on Lasers and Electro-Optics and European Quantum Electronics Conference, ED\_P\_4* (2017), [https://www.osapublishing.org/abstract.cfm?URI=EQEC-2017-ED\\_P\\_4](https://www.osapublishing.org/abstract.cfm?URI=EQEC-2017-ED_P_4).
- [70] R. J. Jones, K. D. Moll, M. J. Thorpe, and J. Ye, Phase-Coherent Frequency Combs in the Vacuum Ultraviolet via High-Harmonic Generation inside a Femtosecond Enhancement Cavity, *Phys. Rev. Lett.* **94**, 193201 (2005).
- [71] A. M. Warrier, J. Lin, H. M. Pask, R. P. Mildren, D. W. Coutts, and D. J. Spence, Highly efficient picosecond diamond Raman laser at 1240 and 1485 nm, *Opt. Express* **22**, 3325 (2014).
- [72] H. K. Raut, V. A. Ganesh, A. S. Nair, and S. Ramakrishna, Anti-reflective coatings: A critical, in-depth review, *Energy Environ. Sci.* **4**, 3779 (2011).
- [73] M. Roth, N. Angert, M. Tseitlin, and A. Alexandrovski, On the optical quality of KTP crystals for nonlinear optical and electro-optic applications, *Opt. Mater.* **16**, 131 (2001).
- [74] M. Leidinger, K. Buse, and I. Breunig, Highly sensitive absorption measurements in lithium niobate using whispering gallery resonators, *Proc. SPIE* **9347**, 93471D (2015).
- [75] N. Waasem, S. Fieberg, J. Hauser, G. Gomes, D. Haertle, F. Kühnemann, and K. Buse, Photoacoustic absorption spectrometer for highly transparent dielectrics with parts-per-million sensitivity, *Rev. Sci. Instrum.* **84**, 023109 (2013).
- [76] A. Yariv and P. Yeh, *Optical Waves in Crystals, Propagation and Control of Laser Radiation* (John Wiley and Sons, New York, 1984).
- [77] X. Luo, L. Wu, J. Chen, Q. Guan, K. Gao, Z.-F. Xu, L. You, and R. Wang, Tunable atomic spin-orbit coupling synthesized with a modulating gradient magnetic field, *Sci. Rep.* **6**, 18983 (2016).
- [78] P. G. Drazin, *Nonlinear Systems* (Cambridge University Press, Cambridge, England, 1992).
- [79] C. Chin, R. Grimm, P. Julienne, and E. Tiesinga, Feshbach resonances in ultracold gases, *Rev. Mod. Phys.* **82**, 1225 (2010).
- [80] W. Zheng, B. Liu, J. Miao, C. Chin, and H. Zhai, Strong Interaction Effects and Criticality of Bosons in Shaken Optical Lattices, *Phys. Rev. Lett.* **113**, 155303 (2014).
- [81] T. Bilitewski and N. R. Cooper, Scattering theory for Floquet-Bloch states, *Phys. Rev. A* **91**, 033601 (2015).
- [82] B. Nagorny, Th. Elsässer, and A. Hemmerich, Collective Atomic Motion in an Optical Lattice Formed inside a High Finesse Cavity, *Phys. Rev. Lett.* **91**, 153003 (2003).
- [83] H. Walther, Experiments on cavity quantum electrodynamics, *Phys. Rep.* **219**, 263 (1992).
- [84] M. Keller, B. Lange, K. Hayasaka, W. Lange, and H. Walther, Continuous generation of single photons with controlled waveform in an ion-trap cavity system, *Nature (London)* **431**, 1075 (2004).
- [85] A. Wallraff, D. I. Schuster, A. Blais, L. Frunzio, R.-S. Huang, J. Majer, S. Kumar, S. M. Girvin, and R. J. Schoelkopf, Strong coupling of a single photon to a superconducting qubit using circuit quantum electrodynamics, *Nature (London)* **431**, 162 (2004).
- [86] C. M. Wilson, G. Johansson, A. Pourkabirian, M. Simoen, J. R. Johansson, T. Duty, F. Nori, and P. Delsing, Observation of the dynamical Casimir effect in a superconducting circuit, *Nature (London)* **479**, 376 (2011).
- [87] F. Piazza and H. Ritsch, Self-Ordered Limit Cycles, Chaos, and Phase Slippage with a Superfluid inside an Optical Resonator, *Phys. Rev. Lett.* **115**, 163601 (2015).
- [88] Y. Chen, Z. Yu, and H. Zhai, Superradiance of Degenerate Fermi Gases in a Cavity, *Phys. Rev. Lett.* **112**, 143004 (2014).
- [89] J.-S. Pan, X.-J. Liu, W. Zhang, W. Yi, and G.-C. Guo, Topological Superradiant States in a Degenerate Fermi Gas, *Phys. Rev. Lett.* **115**, 045303 (2015).