*Moller (Ed.)**Progress in Brain Research*, Vol. 157

ISSN 0079-6123

Copyright © 2006 Elsevier B.V. All rights reserved

CHAPTER 7

Cortical plasticity and rehabilitation

Raluca Moucha and Michael P. Kilgard**Neuroscience Program, School of Brain and Behavioral Sciences, University of Texas at Dallas, TX, USA*

Abstract: The brain is constantly adapting to environmental and endogenous changes (including injury) that occur at every stage of life. The mechanisms that regulate neural plasticity have been refined over millions of years. Motivation and sensory experience directly shape the rewiring that makes learning and neurological recovery possible. Guiding neural reorganization in a manner that facilitates recovery of function is a primary goal of neurological rehabilitation. As the rules that govern neural plasticity become better understood, it will be possible to manipulate the sensory and motor experience of patients to induce specific forms of plasticity. This review summarizes our current knowledge regarding factors that regulate cortical plasticity, illustrates specific forms of reorganization induced by control of each factor, and suggests how to exploit these factors for clinical benefit.

Keywords: Cortical plasticity; Experience-dependent plasticity; Cortical reorganization; Neuromodulators; Cholinergic; Rehabilitation

Factors that regulate plasticity

Plasticity is the remarkable ability of developing, adult, and aging brains to adapt to a changing world. This potential is revealed whenever an organism must meet a new environmental demand or recover from nervous system damage. Plasticity occurs in sensory and motor systems following deprivation of input or overstimulation, increased or decreased usage, learning of new skills, and injury. These experience-dependent changes can be as subtle as a change in neuronal excitability (Engineer et al., 2004) or as dramatic as the rewiring of auditory cortex to process visual information (Sur et al., 1988). Topographic maps, receptive field (RF) size, neuronal firing rate, temporal precision, and combination sensitivity can all be modified by our experiences. The types of plasticity activated by specific situations depend on the nature of the experiences and their behavioral sig-

nificance, conveyed by release of modulatory neurotransmitters (Fig. 1).

Attentional modulation

Neural plasticity is essential for adapting to changes in the environment but plasticity can be destabilizing if not well regulated. Limiting plasticity prevents meaningless events from driving changes that could degrade previously acquired memories and skills. Attention plays a key role in the regulation of plasticity associated with sensory experience. Repeated sensory stimulation alters topography in primary sensory cortex only when monkeys use the stimuli to make behavioral judgments (Recanzone et al., 1992, 1993). Many studies have shown that cortical neurons respond differently to attended versus unattended stimuli. Neurons in secondary somatosensory cortex, for example, exhibit greater response synchronization when monkeys are engaged in a tactile task (Steinmetz et al., 2000). Attention can also directly affect

*Corresponding author. Tel.: (972) 883-2345; Fax: (972) 883-2491; E-mail: kilgard@utdallas.edu

Fig. 1. Several neurotransmitter systems which project widely into the cortex are implicated in learning and experience-dependent plasticity: Acetylcholine from the cholinergic nucleus basalis (NB), dopamine from the ventral tegmentum (VTA), noradrenaline from the locus coeruleus (LC), and serotonin from the raphe nuclei (RN). In addition to these major neurotransmitters, GABA-ergic projections, histamine, and neuro-hormones also play a role in modulating plasticity. Release of these transmitters is normally regulated by behavioral state but can also be triggered by drugs or direct electrical stimulation. Cortical plasticity results when release of these transmitters is repeatedly associated with the occurrence of a sensory stimulus. (Source: Figure adapted from McEwen BS, 2003, Karger Gazette, No. 66, Basel, S. Karger AG.)

firing rates of cortical neurons (Treue and Maunsell, 1999; Recanzone and Wurtz, 2000). Results from several psychophysical studies support the hypothesis that attention regulates cortical plasticity and learning. Distinct forms of perceptual learning result when subjects attend to different features of an otherwise identical sensory input (Ahissar and Hochstein, 1993). Exposure to moving dot patterns can improve motion direction discrimination ability even if the motion is undetectable (due to low coherence), as long as the subjects are actively engaged in a visual task (Watanabe et al., 2001; Seitz and Watanabe, 2003). These results suggest that directed attention facilitates the learning of associated sensory features.

Neuromodulatory influences

Several neuromodulators, such as dopamine, nor-epinephrine, and acetylcholine, are known to regulate learning and memory in humans (Hasselmo, 1995). The observation that synaptic plasticity is also enhanced by the presence of these neurotransmitters supports the relationship between learning and plasticity (Singer, 1986; Brocher et al., 1992). Injection of acetylcholine or nor-epinephrine directly into visual, somatosensory, or auditory cortex during sensory stimulation can promote expression of neural plasticity in the intact brain (Greuel et al., 1988; McKenna et al., 1989; Delacour et al., 1990). Pairing sensory inputs with electrical activation of the nucleus basalis (NB), locus coeruleus (LC), or ventral tegmental area (VTA) also results in plasticity that is specific to features of the associated input (Kilgard and

Merzenich, 1998; Bouret and Sara, 2002; Bao et al., 2003). Stimulation of neuromodulatory neurotransmitter release by amphetamine enhances cortical plasticity in human subjects (Dinse et al., 2003; Tegenthoff et al., 2004). Since release of these neurotransmitters is normally triggered by behaviorally arousing events, it is likely they contribute to the regulation of cortical plasticity.

Patterns of sensory activation

Many studies have shown that sensory input determines the form of cortical reorganization. When animals or humans repeatedly practice a skill that engages a limited region of the sensory epithelium, the regions of the cortical map that respond to task-specific inputs are enlarged (Jenkins et al., 1990; Recanzone et al., 1992, 1993; Elbert et al., 1995; Sterr et al., 1998). Cortical RFs can narrow or broaden and response latency can increase or decrease depending on the spatial and temporal pattern of sensory activation encountered during training. Owl monkeys trained on a tone frequency discrimination task have A1 neurons with smaller RFs and longer response latencies than untrained controls (Recanzone et al., 1993). Monkeys trained to detect changes in the rate of a tactile vibration exhibit larger RFs and faster response latencies (Recanzone et al., 1992). In contrast, training on a task with stimuli that move across the skin cause RFs to shrink (Jenkins et al., 1990). Training on a visual orientation task increased the steepness of orientation tuning in the trained region of the visual field (Schoups et al., 2001). These studies support the hypothesis that perceptual learning and cortical plasticity are specific to attended sensory features.

The rodent whisker system has proven particularly useful for directly comparing how cortical plasticity is shaped by different spatial patterns of activity. If all but one whisker is cut, for example, the responsiveness of the spared whisker is increased (Glazewski et al., 1998). Cutting a single whisker reduces input to the corresponding region of barrel cortex, decreases the responsiveness of the deprived neurons, and increases the responsiveness to neighboring whiskers. If all the whisk-

ers are cut, the reduction in the response to the principle whisker is more modest. A checkerboard deprivation pattern causes responses to the deprived whiskers to decrease, but does not increase the response to the spared whiskers (Wallace and Fox, 1999). Finally, cutting all but two neighboring whiskers causes the RF of neurons in each region to shift toward the other spared whisker (Diamond et al., 1993). These results suggest that competition between sensory inputs induces the different forms of changes in responsiveness.

Timing of sensory inputs

The temporal coincidence of sensory stimulation can be just as important as its spatial pattern in determining the direction and magnitude of cortical plasticity. Inputs that are correlated in time are more likely to cause a change in neural responses than uncorrelated inputs. Simultaneous activation of an area of skin with a vibrating disc increased RF size in primary somatosensory cortex, while stimulation of a single point on the skin does not cause any change (Godde et al., 1996). Simultaneous activation of the developing auditory system by repeated exposure to broadband noise causes increased cortical RFs and degraded tonotopic maps (Chang and Merzenich, 2003). Such changes are not seen after equivalent exposure to tones. Increased simultaneous activation of the fingers due to surgical fusion or operant training leads to large, multidigit RFs in somatosensory cortex (Allard et al., 1991; Wang et al., 1995). This finding suggests that the usual segregation of each digit's cortical representation reflects the normally asynchronous activation of each digit. In vitro and more recently in vivo studies have further demonstrated that the time window for correlated inputs to induce plasticity is on the order of tens of milliseconds (Tsodyks, 2002; Dan and Poo, 2004). These results indicate that the precise spatial and temporal pattern of inputs shape cortical networks due to operation of Hebbian synaptic plasticity.

1 *Duration of experience*

3 Many factors regulate the time course of learning
 5 and plasticity (Ebbinghaus, 1885; Dubnau et al.,
 7 2003). Fear conditioning can induce rapid and
 9 long lasting shifts of neuronal tuning toward the
 11 frequency of the conditioned tone (Bjordahl et al.,
 13 1998; Weinberger, 2003). In contrast, plasticity
 15 following skill learning or use-dependent plasticity
 17 develops gradually over time. The magnitude of
 19 effects often depends on duration of training and
 21 correlates with performance accuracy (Pleger et
 23 al., 2003). Motor map reorganization, which is
 25 accompanied by synaptogenesis and believed to
 27 underlie consolidation of motor skills, occurs dur-
 29 ing the late phase (after 10 days) of motor skill
 31 learning (Kleim et al., 2004). The schedule of in-
 33 puts can also determine the induction of stable
 35 versus reversible synaptic modifications (Ma-
 37 uelshagen et al., 1998). Spaced repetition of LTP
 39 inducing stimuli prevents the reversal of LTP due
 41 to subsequent spontaneous activity that occurs af-
 43 ter massed repetition (Zhou et al., 2003). Stable
 45 synaptic modifications are also induced by visual
 47 experience when the exposure to unidirectional
 moving bars occurs in a spaced pattern (three sets
 of 60 flashes separated by 5 min) versus massed
 pattern (180 flashes continuously). If persistent
 synaptic changes are important for learning and
 memory, the effective use of training strategies
 that prevent their reversal is important. In behav-
 iorally trained mice temporally spaced training
 more effectively recruits protein synthesis and en-
 hanced long-term memory of contextual condi-
 tioning, while massed training triggers greater
 protein phosphatase 1 activity which suppresses
 memory formation (Genoux et al., 2002; Scharf et
 al., 2002). These results suggest that the schedule
 of training determines the duration of neural plas-
 ticity and learning.

43 **Influence of background stimuli on plasticity**

45 Psychologists and psychophysicists have known
 47 for decades that unattended background stimuli
 (context) influence perceptual learning. Studies of
 sensory plasticity have typically been conducted in

environments stripped of context, by using sound-
 proof booths or gray backgrounds. Recent exper-
 iments in more naturalistic and complex settings
 have shown that context also influences plasticity.

In many cases, adding complex backgrounds
 actually improves learning. Contrast discrimina-
 tion learning, for example, can be facilitated by
 fixed contrast stimuli flanking the target stimulus
 (Adini et al., 2002). Dim line objects are easier to
 detect when flanked with a second collinear bar
 (Kapadia et al., 1995). This improvement in hu-
 man performance is paralleled an enhancement of
 neuronal responses in monkey V1 when equivalent
 visual stimuli are presented (Kapadia et al., 1995).
 When foot shock is paired with a tone, the pres-
 ence of unpaired background tones determines
 whether auditory cortex neurons shift their fre-
 quency tuning toward or away from the paired
 tone (Bakin and Weinberger, 1990; Ohl and
 Scheich, 1996; Dimyan and Weinberger, 1999).

While all of these studies suggest that many
 factors regulate plasticity and learning, direct
 comparison of the interactions between these fac-
 tors has proven difficult. Differences in the be-
 havioral response, task difficulty, task goal,
 motivation, modality and species often confound
 the influence of the discussed factors on plasticity.
 Because these factors are so tightly interdependent
 it has been difficult to tease apart their relative
 importance in directing different forms of plastic-
 ity. Varying sensory patterns or adding a complex
 background, for example, would also affect task
 difficulty in most cases. Currently, reduced prep-
 arations provide the best opportunities to study
 the interactions between each of the factors that
 regulate neural plasticity. Experimental paradigms
 that directly stimulate modulatory systems have
 proven particularly valuable in documenting the
 influence of stimulus pattern, timing, and back-
 ground conditions on cortical plasticity.

Sensory input paired with controlled release of neuromodulators

Pairing electrical activation of the cholinergic NB
 with different sounds generates changes in cortical
 map and RF properties in rats that closely parallel

the different forms of plasticity resulting from operant training in monkeys. For example, temporally modulated stimuli tend to *increase* RF size, while stimuli that activate different regions of the receptor surface tend to *decrease* RF size (Kilgard et al., 2002). While the differential plasticity observed in operant studies could be attributed to any number of technical differences, in the NB stimulation experiments the only explanation for the differential plasticity was the temporal and spectral properties of the sounds associated with NB stimulation. The observation that similar sensory inputs lead to comparable plasticity even in the absence of operant training supports the conclusion that sensory features determine the form of cortical plasticity.

During natural learning, changing task contingencies are known to alter the type, amount, and timing of neuromodulator release. For example, novel sounds activate cholinergic NB neurons for a few trials, but habituate rapidly (Richardson and DeLong, 1990, 1991). The response can later be reinstated if the sound is associated with a reward or punishment. NB releases acetylcholine onto the cortex only during the learning phase of a lever press task, but not after the task is well learned (Orsetti et al., 1996). Electrical stimulation bypasses the natural triggers of NB activity and eliminates the natural brake on cortical plasticity. The consistency of electrical activation makes it possible to systematically compare how the type, amount, and timing of neuromodulator release influence cortical plasticity when associated with sensory stimuli of differing spatial and temporal properties.

Patterns of activation determine type of reorganization

Distinct types of cortical reorganization are generated when NB stimulation is associated with different sensory inputs. Cortical topography, RF size, and response timing are altered as a function of the temporal modulation and spatial distribution of inputs associated with NB stimulation. The focal activation caused by presentations of a single tone frequency results in expansion of the area responsive to the tone, and modest RFs broadening. Distributing the activation over more frequency sectors (i.e., seven tone frequencies) prevents the map reorganization but results in a narrowing of RFs (Kilgard et al., 2001). Rapidly modulated tone trains cause map expansion and dramatic RF broadening when activation is focal (i.e., one carrier frequency) and less extreme RF broadening and no map plasticity when the tone trains activate several regions (i.e., seven different carrier frequencies). These results document how different activation patterns direct cortical plasticity: (1) sensory map expansion only results when sensory activation is focal. (2) Distributing inputs across the cochlea tends to reduce RF size. (3) Modulated stimuli tend to increase RF size compared to unmodulated stimuli (Table 1, Kilgard et al., 2002). The observation that RF size is increased by stimuli with high degree of temporal modulation and little spatial variability (tone trains) and decreased by stimuli with high spatial variability and no temporal modulation (unmodulated tones of varying frequency) is consistent with earlier observations of plasticity in operant trained monkeys (Recanzone et al., 1992, 1993). These results indicate that NB stimulation directs changes that are

Table 1. Plasticity induced by pairing NB with different sounds

NB stimulation paired with	Plasticity observed	References
Single tone	Map expansion + decrease latency	Kilgard and Merzenich (1998)
Tone train	Map expansion + decrease latency + RF broadening	Kilgard and Merzenich, 2001
Distributed tones	RF narrowing + increase latency	Kilgard and Merzenich (1998)
Distributed tone trains	RF broadening + temporal plasticity	Kilgard and Merzenich (2001)
Frequency modulated tones	RF broadening + decrease latency + decreased thresholds	Moucha et al. (2005)
Complex acoustic sequence	Combination sensitivity + decrease latency + decreased thresholds	Kilgard and Merzenich (2002)
Background sounds	Alters plasticity generated in silence	Moucha et al. (2005)

similar to operant induced plasticity even though the rats did not use the stimuli in any way.

Natural sounds usually vary both in spatial and temporal structure and create more complex activity patterns than tones. Pairing frequency modulated sweeps and complex acoustic sequences leads to forms of plasticity that are unpredictable from earlier studies with simple tones (Kilgard and Merzenich, 2002; Moucha et al., 2005). FM sweeps result in a moving pattern of activation across the cochlea. Although tones and FM sweeps may share many features in common (including starting frequency, bandwidth, intensity, duration), pairing FM sweeps with NB stimulation causes different plasticity compared with unmodulated tones. Pairing FM sweeps with NB decreases response latency, broadens RFs, and increased sensitivity to quiet tones. These changes are restricted to the region of A1 activated by the sweep, but no map expansion results. When the starting frequency of the FM sweeps is varied no plasticity is observed in any region of A1 (Moucha et al., 2005). While repeated exposure to FM's does not cause any preference for FM direction (increasing or decreasing), pairing a sequence of sounds with identical NB stimulation can result in the development of responses sensitive to tone order.

Although plasticity mechanisms have presumably evolved to increase cortical processing capacity for behaviorally relevant inputs, it is not immediately obvious why the plasticity associated with each spatial and temporal input pattern is beneficial.

Correlation of sensory inputs

Studies in auditory, visual, and somatosensory cortex have suggested that input correlations strongly influence neural plasticity (Buonomano and Merzenich, 1998). In the developing visual system, for example, alternating asynchronous electrical stimulation of the optic nerve prevents normal development of binocular visual responses (Stryker and Strickland, 1984). In auditory cortex, sounds designed to decrease or increase correlation across the frequency map lead to very different forms of plasticity (Pandya et al., 2005). Alternat-

ing activation of two nonoverlapping auditory neuron populations by two tones of distant frequencies (2 and 14 kHz) results in map segregation, decreased excitability, and longer response latencies of the activated neurons. These changes do not occur when NB-stimulation is paired with a modulated noise burst that synchronously activate large populations of A1 neurons. Pairing pulsed noises with NB stimulation disrupts tonotopic maps and reduces spontaneous discharge correlation in the primary auditory cortex (Bao et al., 2003). These findings are in agreement with the Hebbian postulate that inputs with decreased correlation weaken cortical responses and supports other observations that primary sensory cortices segregate inputs that are asynchronous and integrate correlated inputs (Allard et al., 1991; Wang et al., 1995).

Duration of associative sensory pairing

The duration of NB-induced plasticity depends on the schedule of the pairing protocol. Repetitively pairing NB stimulation with a tone for several minutes causes a shift in frequency tuning that reverses within 5 h (Zhang et al., 2005). Cortical map expansion builds with repeated pairings. One month of 300 NB-tone pairings per day increases the A1 representation of the paired frequency by twice as much as a week of pairing (Kilgard and Merzenich, 1998). After a month of pairing, NB-induced map plasticity endures for at least 20 days (Carrasco et al., 2004). NB stimulation also increases the duration of cortical and subcortical plasticity induced by cortical microstimulation (Ma and Suga, 2003). These results support earlier observations that cholinergic modulation contributes to both short-term and long-term plasticity.

Background stimuli influence plasticity outcomes

Although background stimuli are known to influence task performance and plasticity (Kapadia et al., 1995; Adini et al., 2002), it has not been clear whether the differences are due to altered task difficulty or to some specific influence of the dis-

tracters. By directly pairing sensory stimuli with NB stimulation in different contexts, we have shown that background stimuli can influence plasticity independent of any influence on task performance.

Background sounds can alter bandwidth, threshold, and map plasticity. The 20% increase in RF size that occurs after pairing a single tone with NB stimulation does not occur if the same tone-NB pairing is interleaved with flanking tones that are not associated with NB stimulation (Kilgard et al., 2001). Repeated presentation of the word /SASH/ paired with NB stimulation causes expansion of the high frequency region of A1 (Pandya et al., 2003), presumably because the first element of the word is the high frequency /S/ sound. This map plasticity is eliminated when each phoneme of the word /S/, /A/, and /SH/ are also presented, but not paired with NB stimulation. Finally, the addition of unpaired FM sweeps that contrast the duration and direction of the paired FM sweeps results in threshold and latency plasticity not observed if the identical FM's sweeps are paired with NB stimulation in a silent background (Moucha et al., 2005). These results indicate that background conditions, previously thought to be irrelevant, are likely to shape many forms of cortical plasticity.

Clinical conclusions

It was proposed two decades ago that cortical reorganization after injury may be the neural substrate for recovery of function after brain damage (Jenkins and Merzenich, 1987). More recent studies in primates have shown that rehabilitative training can direct reorganization to benefit recovery (Nudo et al., 1996). There is no longer a doubt that reorganization after brain lesions is shaped by the sensorimotor experiences in the weeks to months following injury. Hence it is important to effectively manage plasticity after brain damage. Many of the factors that influence plasticity can be manipulated in clinical settings to enhance therapeutic outcomes.

Attention is often impaired after brain injury and likely plays a critical role in directing training-

induced plasticity. Patients with the highest vigilance scores typically receive greatest benefit from the rehabilitation therapy (Sohlberg et al., 2000). Some strategies, such as constraint therapy, that increase arousal (and even frustration) can be more effective than traditional occupational therapies (Taub and Uswatte, 2003).

The diffuse modulatory systems including the cholinergic NB are particularly vulnerable to dysregulation after brain damage. Experimental damage to the NB prevents map reorganization and retards skill learning in rats (Fig. 2). The recent observation that NB damage also prevents recovery from brain damage suggests many of the same mechanisms that regulate normal learning also regulate recovery from injury (Conner et al., 2005). In some patients, medication may be beneficial for normalizing attentional and neuromodulatory mechanisms. Agents that stimulate neuromodulators known to place the brain in a permissive state for experience-dependent changes are most likely to be effective.

Drugs that act on noradrenergic, dopaminergic, serotonergic, and cholinergic systems have been shown in laboratory and clinical research to be pharmacological adjuvants in neurorehabilitation (Phillips et al., 2003). Amphetamines lead to a diffuse increase of several modulators and can have a positive influence even when administered only as a single dose at the beginning of therapy (Feeney et al., 1982). It is important to note that drug administration only aids recovery when paired with practice. Amphetamine administration during speech language therapy increases the rate of improvement of aphasic patients during the early recovery period after stroke (Walker-Batson et al., 2004). Amphetamine also facilitates speech training in adult cochlear implant users (Tobey et al., 2005) and second language acquisition in normal subjects (Breitenstein et al., 2004). More research is needed to evaluate how best to facilitate neurological recovery using nervous system stimulants and other psychoactive compounds.

Since sensory and motor experiences (associated with release of modulatory neurotransmitters) determine the form of plasticity generated, it is critical to develop targeted rehabilitation techniques designed to stimulate adaptive plasticity following

Fig. 2. Rats can improve their grasping skill with repeated practice. After practice more neurons in caudal motor cortex control the trained forepaw (b). Without the normal input from the cholinergic nucleus basalis (i.e., after lesion of NB cholinergic neurons) rats cannot improve their accuracy with training and the motor map of the trained forepaw remains unchanged (c). This result indicates that practice alone without appropriate levels of neuromodulators does not result in learning or map plasticity. Cortical lesions of the caudal forepaw representation after training (d) results in loss of accuracy that can be recovered after retraining and expansion of the rostral motor map of the trained forepaw (e). If nucleus basalis is lesioned during retraining, recovery of reaching accuracy is impaired and the rostral forepaw representation does not change (f). This result indicates that appropriate levels of neuromodulators are also required to promote compensatory plasticity and recovery of function after brain damage. (Source: Results illustrated are from experiments by Conner et al., 2005).

brain damage. Motor maps are altered by skill acquisition not by repetitive use alone (Nudo, 1997). In somatosensory cortex postlesion changes are related to individual strategies and sensorimotor experience resulting from idiosyncratic behavior. The type of reorganization often depends on the strategy used by individual monkeys to re-acquire an object retrieval skill after an experimentally-induced stroke (Xerri et al., 1998). These findings imply that cortical map plasticity can be influenced by the pattern of sensorimotor stimulation during behavioral treatment. In dysphagic stroke patients electrical stimulation of the phar-

ynx results in motor cortex plasticity that is dependent on the pattern of stimulation (frequency, intensity, and duration of stimulation) and correlates with improvement in swallowing function (Fraser et al., 2002). Several treatment strategies now effectively combine modulation of somatosensory input, administration of pharmacological adjuvants, and cortical stimulation to improve outcomes of rehabilitation (Hummel and Cohen, 2005).

The influence of background has not been well-studied in the context of neurorehabilitation. However, studies have documented beneficial

Table 2. Factors regulating plasticity

Factors regulating plasticity	Effect	References
Attention	Enhances stimulus driven plasticity via internal trigger of neuromodulator release	Hasselmo (1995)
Drugs	Achieve optimal levels of neuromodulators required for plasticity	Phillips et al. (2003)
Pattern of stimuli	Determines form of plasticity (reorganization of sensory representations, temporal precision, spatial selectivity, etc.)	Buonomano and Merzenich (1998)
Temporal delivery (spaced vs. massed training)	Potentiation and stabilization of changes by stimulating protein synthesis mechanisms, and reducing phosphatases that prevent long-term changes	Genoux et al. (2002); Scharf et al. (2002); Zhou et al. (2003)
Duration of training	Consolidation of changes via synaptogenesis	Klein et al. (2004)

effects of general environmental enrichment in recovery after experimental brain infarcts (review Johansson, 2004). Enriched environments further enhance recovery when combined with training or drug therapy (Biernaskie and Corbett, 2001; Puurunen et al., 2001). Our results from plasticity experiments indicate that adding complex backgrounds during rehabilitative training may aid in emphasizing and facilitating performance on specific tasks.

In conclusion, therapies that optimize neural plasticity by integrating all the concepts described above are likely to improve patient outcomes (Table 2). Optimal modulator release can be accomplished by modulating attention and arousal either through task requirements or stimulating drugs. Stimuli used in training can be selected to address specific changes (rewiring) needed to direct recovery of function in individual patients. The proper timing of training sessions (i.e., spaced rather than massed training) and duration should also be optimized for training to be effective and long lasting. The addition of background stimuli may prove beneficial in many situations. This context can be used to emphasize aspects of a task or to incrementally increase task difficulty to maintain the patient's motivation and arousal. Ideally, the progress and efficacy of therapy should be monitored (and adjusted) in each patient using brain imaging or evoked potentials.

We are now beginning to understand how many factors interplay in directing different forms of plasticity. Manipulation of the many parameters known to shape brain plasticity, including the pattern, timing, and duration of events associated

with attention and release of modulatory neurotransmitters, is essential to improving neurorehabilitation.

Acknowledgments

We thank Cherie Percaccio, Amanda Puckett, Vikram Jakkamsetti, Dr. Owen Floddy and Dr. Pritesh Pandya for insightful comments and review of the manuscript. This work was supported by grants from the National Institute for Deafness and Other Communication Disorders, the James S. McDonnell Foundation, and Cure Autism Now.

References

- Adini, Y., Sagi, D. and Tsodyks, M. (2002) Context-enabled learning in the human visual system. *Nature*, 415: 790–793.
- Ahissar, M. and Hochstein, S. (1993) Attentional control of early perceptual learning. *Proc. Natl. Acad. Sci. USA.*, 90: 5718–5722.
- Allard, T., Clark, S.A., Jenkins, W.M. and Merzenich, M.M. (1991) Reorganization of somatosensory area 3b representations in adult owl monkeys after digital syndactyly. *J. Neurophysiol.*, 66: 1048–1058.
- Bakin, J.S. and Weinberger, N.M. (1990) Classical conditioning induces CS-specific receptive field plasticity in the auditory cortex of the guinea pig. *Brain Res.*, 536: 271–286.
- Bao, S., Chang, E.F., Davis, J.D., Gobeske, K.T. and Merzenich, M.M. (2003) Progressive degradation and subsequent refinement of acoustic representations in the adult auditory cortex. *J. Neurosci.*, 23: 10765–10775.
- Biernaskie, J. and Corbett, D. (2001) Enriched rehabilitative training promotes improved forelimb motor function and enhanced dendritic growth after focal ischemic injury. *J. Neurosci.*, 21: 5272–5280.

- 1 Bjordahl, T.S., Dimyan, M.A. and Weinberger, N.M. (1998) Induction of long-term receptive field plasticity in the auditory cortex of the waking guinea pig by stimulation of the nucleus basalis. *Behav. Neurosci.*, 112: 467–479.
- 3 Bouret, S. and Sara, S.J. (2002) Locus coeruleus activation modulates firing rate and temporal organization of odour-induced single-cell responses in rat piriform cortex. *Eur. J. Neurosci.*, 16: 2371–2382.
- 5 Breitenstein, C., Wailke, S., Bushuven, S., Kamping, S., Zwitterlood, P., Ringelstein, E.B. and Knecht, S. (2004) D-amphetamine boosts language learning independent of its cardiovascular and motor arousing effects. *Neuropsychopharmacology*, 29: 1704–1714.
- 7 Brocher, S., Artola, A. and Singer, W. (1992) Agonists of cholinergic and noradrenergic receptors facilitate synergistically the induction of long-term potentiation in slices of rat visual cortex. *Brain Res.*, 573: 27–36.
- 9 Buonomano, D.V. and Merzenich, M.M. (1998) Cortical plasticity: from synapses to maps. *Annu. Rev. Neurosci.*, 21: 149–186.
- 11 Carrasco, R.A., Puckett, A., Pandya, P.K., McMenamy, A., Gibbons, J., Heydrick, C., Moucha, R., Jain, R. and Kilgard, M.P. (2004). Frequency map reorganization in rat primary auditory cortex enabled by basal forebrain activation endures for at least 20 days. In: Society for Neuroscience, San Diego, CA.
- 13 Chang, E.F. and Merzenich, M.M. (2003) Environmental noise retards auditory cortical development. *Science*, 300: 498–502.
- 15 Conner, J.M., Chiba, A.A. and Tuszynski, M.H. (2005) The basal forebrain cholinergic system is essential for cortical plasticity and functional recovery following brain injury. *Neuron*, 46: 173–179.
- 17 Dan, Y. and Poo, M.M. (2004) Spike timing-dependent plasticity of neural circuits. *Neuron*, 44: 23–30.
- 19 Delacour, J., Houcine, O. and Costa, J.C. (1990) Evidence for a cholinergic mechanism of “learned” changes in the responses of barrel field neurons of the awake and undrugged rat. *Neuroscience*, 34: 1–8.
- 21 Diamond, M.E., Armstrong-James, M. and Ebner, F.F. (1993) Experience-dependent plasticity in adult rat barrel cortex. *Proc. Natl. Acad. Sci. USA*, 90: 2082–2086.
- 23 Dimyan, M.A. and Weinberger, N.M. (1999) Basal forebrain stimulation induces discriminative receptive field plasticity in the auditory cortex. *Behav. Neurosci.*, 113: 691–702.
- 25 Dinse, H.R., Ragert, P., Pleger, B., Schwenkreis, P. and Tegenthoff, M. (2003) Pharmacological modulation of perceptual learning and associated cortical reorganization. *Science*, 301: 91–94.
- 27 Dubnau, J., Chiang, A.S. and Tully, T. (2003) Neural substrates of memory: from synapse to system. *J. Neurobiol.*, 54: 238–253.
- 29 Ebbinghaus, H. (1885) *Über das Gedächtnis. Untersuchungen zur Experimentellen Psychologie*. Duckner & Humboldt, Leipzig.
- 31 Elbert, T., Pantev, C., Wienbruch, C., Rockstroh, B. and Taub, E. (1995) Increased cortical representation of the fingers of the left hand in string players. *Science*, 270: 305–307.
- 33 Engineer, N.D., Percaccio, C.R., Pandya, P.K., Moucha, R., Rathbun, D.L. and Kilgard, M.P. (2004) Environmental enrichment improves response strength, threshold, selectivity, and latency of auditory cortex neurons. *J. Neurophysiol.*, 92: 73–82.
- 35 Feeney, D.M., Gonzalez, A. and Law, W.A. (1982) Amphetamine, haloperidol, and experience interact to affect rate of recovery after motor cortex injury. *Science*, 217: 855–857.
- 37 Fraser, C., Power, M., Hamdy, S., Rothwell, J., Hobday, D., Hollander, I., Tyrell, P., Hobson, A., Williams, S. and Thompson, D. (2002) Driving plasticity in human adult motor cortex is associated with improved motor function after brain injury. *Neuron*, 34: 831–840.
- 39 Genoux, D., Haditsch, U., Knobloch, M., Michalon, A., Storm, D. and Mansuy, I.M. (2002) Protein phosphatase 1 is a molecular constraint on learning and memory. *Nature*, 418: 970–975.
- 41 Glazewski, S., McKenna, M., Jacquin, M. and Fox, K. (1998) Experience-dependent depression of vibrissae responses in adolescent rat barrel cortex. *Eur. J. Neurosci.*, 10: 2107–2116.
- 43 Godde, B., Spengler, F. and Dinse, H.R. (1996) Associative pairing of tactile stimulation induces somatosensory cortical reorganization in rats and humans. *Neuroreport.*, 8: 281–285.
- 45 Greuel, J.M., Luhmann, H.J. and Singer, W. (1988) Pharmacological induction of use-dependent receptive field modifications in the visual cortex. *Science*, 242: 74–77.
- 47 Hasselmo, M.E. (1995) Neuromodulation and cortical function: modeling the physiological basis of behavior. *Behav. Brain Res.*, 67: 1–27.
- Hummel, F.C. and Cohen, L.G. (2005) Drivers of brain plasticity. *Curr. Opin. Neurol.*, 18: 667–674.
- Jenkins, W.M. and Merzenich, M.M. (1987) Reorganization of neocortical representations after brain injury: a neurophysiological model of the bases of recovery from stroke. *Prog. Brain Res.*, 71: 249–266.
- Jenkins, W.M., Merzenich, M.M., Ochs, M.T., Allard, T. and Guic-Robles, E. (1990) Functional reorganization of primary somatosensory cortex in adult owl monkeys after behaviorally controlled tactile stimulation. *J. Neurophysiol.*, 63: 82–104.
- Johansson, B.B. (2004) Brain plasticity in health and disease. *Keio J. Med.*, 53: 231–246.
- Kapadia, M.K., Ito, M., Gilbert, C.D. and Westheimer, G. (1995) Improvement in visual sensitivity by changes in local context: parallel studies in human observers and in V1 of alert monkeys. *Neuron*, 15: 843–856.
- Kilgard, M.P. and Merzenich, M.M. (1998) Cortical map reorganization enabled by nucleus basalis activity. *Science*, 279: 1714–1718.
- Kilgard, M.P. and Merzenich, M.M. (2002) Order sensitive plasticity in adult primary auditory cortex. *Proc. Natl. Acad. Sci. USA*, 99: 3205–3209.
- Kilgard, M.P., Pandya, P.K., Engineer, N.D. and Moucha, R. (2002) Cortical network reorganization guided by sensory input features. *Biol. Cybern.*, 87: 333–343.
- Kilgard, M.P., Pandya, P.K., Vazquez, J., Gehi, A., Schreiner, C.E. and Merzenich, M.M. (2001) Sensory input directs spa-

- 1 tial and temporal plasticity in primary auditory cortex. *J. Neurophysiol.*, 86: 326–338.
- 3 Kleim, J.A., Hogg, T.M., VandenBerg, P.M., Cooper, N.R.,
Bruneau, R. and Remple, M. (2004) Cortical synaptogenesis
5 and motor map reorganization occur during late, but not
early, phase of motor skill learning. *J. Neurosci.*, 24: 628–633.
- 7 Ma, X. and Suga, N. (2003) Augmentation of plasticity of the
central auditory system by the basal forebrain and/or so-
matosensory cortex. *J. Neurophysiol.*, 89: 90–103.
- 9 Mauelshagen, J., Sherff, C.M. and Carew, T.J. (1998) Differ-
ential induction of long-term synaptic facilitation by spaced
11 and massed applications of serotonin at sensory neuron
synapses of *Aplysia californica*. *Learn. Mem.*, 5: 246–256.
- 13 McKenna, T.M., Ashe, J.H. and Weinberger, N.M. (1989)
Cholinergic modulation of frequency receptive fields in au-
ditory cortex: I. Frequency-specific effects of muscarinic ago-
nists. *Synapse*, 4: 30–43.
- 15 Moucha, R., Pandya, P.K., Engineer, N.D., Rathbun, D.L. and
Kilgard, M.P. (2005) Background sounds contribute to spec-
17 trotemporal plasticity in primary auditory cortex. *Exp. Brain*
Res., 162: 417–427.
- 19 Nudo, R.J. (1997) Remodeling of cortical motor representa-
tions after stroke: implications for recovery from brain dam-
age. *Mol. Psychiatry*, 2: 188–191.
- 21 Nudo, R.J., Wise, B.M., SiFuentes, F. and Milliken, G.W.
(1996) Neural substrates for the effects of rehabilitative
23 training on motor recovery after ischemic infarct. *Science*,
272: 1791–1794.
- 25 Ohl, F.W. and Scheich, H. (1996) Differential frequency con-
ditioning enhances spectral contrast sensitivity of units in
auditory cortex (field AL) of the alert Mongolian gerbil. *Eur.*
27 *J. Neurosci.*, 8: 1001–1017.
- Orsetti, M., Casamenti, F. and Pepeu, G. (1996) Enhanced
acetylcholine release in the hippocampus and cortex during
29 acquisition of an operant behavior. *Brain Res.*, 724: 89–96.
- 31 Pandya, P.K., Engineer, N., Moucha, R., Rathbun, D.L.,
Vazquez, J. and Kilgard, M.P. (2003). Background stimuli
control cortical map expansion in primary auditory cortex.
33 In: International Conference on Auditory Cortex, Ma-
gdeburg, Germany.
- 35 Pandya, P.K., Moucha, R., Engineer, N.D., Rathbun, D.L.,
Vazquez, J. and Kilgard, M.P. (2005) Asynchronous inputs
alter excitability, spike timing, and topography in primary
37 auditory cortex. *Hear. Res.*, 203: 10–20.
- Phillips, J.P., Devier, D.J. and Feeney, D.M. (2003) Rehabil-
39 itation pharmacology: bridging laboratory work to clinical
application. *J. Head Trauma Rehabil.*, 18: 342–356.
- 41 Pleger, B., Foerster, A.F., Ragert, P., Dinse, H.R., Schwenk-
reis, P., Malin, J.P., Nicolas, V. and Tegenthoff, M. (2003)
Functional imaging of perceptual learning in human primary
43 and secondary somatosensory cortex. *Neuron*, 40: 643–653.
- Puurunen, K., Jolkkonen, J., Sirvio, J., Haapalinna, A. and
45 Sivenius, J. (2001) Selegiline combined with enriched-envi-
ronment housing attenuates spatial learning deficits following
focal cerebral ischemia in rats. *Exp. Neurol.*, 167: 348–355.
- 47 Recanzone, G.H., Merzenich, M.M., Jenkins, W.M., Grajski,
K.A. and Dinse, H.R. (1992) Topographic reorganization of
the hand representation in cortical area 3b owl monkeys
trained in a frequency-discrimination task. *J. Neurophysiol.*,
67: 1031–1056.
- Recanzone, G.H., Schreiner, C.E. and Merzenich, M.M. (1993)
Plasticity in the frequency representation of primary auditory
5 cortex following discrimination training in adult owl mon-
keys. *J. Neurosci.*, 13: 87–103.
- Recanzone, G.H. and Wurtz, R.H. (2000) Effects of attention
on MT and MST neuronal activity during pursuit initiation.
7 *J. Neurophysiol.*, 83: 777–790.
- Richardson, R.T. and DeLong, M.R. (1990) Context-depend-
9 ent responses of primate nucleus basalis neurons in a go/no-
go task. *J. Neurosci.*, 10: 2528–2540.
- Richardson, R.T. and DeLong, M.R. (1991) Electrophysiol-
11 ogy studies of the functions of the nucleus basalis in primates.
Adv. Exp. Med. Biol., 295: 233–252.
- Scharf, M.T., Woo, N.H., Lattal, K.M., Young, J.Z., Nguyen,
15 P.V. and Abel, T. (2002) Protein synthesis is required for the
enhancement of long-term potentiation and long-term mem-
17 ory by spaced training. *J. Neurophysiol.*, 87: 2770–2777.
- Schoups, A., Vogels, R., Qian, N. and Orban, G. (2001) Prac-
tising orientation identification improves orientation coding
19 in V1 neurons. *Nature*, 412: 549–553.
- Seitz, A.R. and Watanabe, T. (2003) Psychophysics: is sublim-
21 inal learning really passive? *Nature*, 422: 36.
- Singer, W. (1986) The brain as a self-organizing system. *Eur.*
23 *Arch. Psychiatry Neurol. Sci.*, 236: 4–9.
- Sohlberg, M.M., McLaughlin, K.A., Pavese, A., Heidrich, A.
and Posner, M.I. (2000) Evaluation of attention process
25 training and brain injury education in persons with acquired
brain injury. *J. Clin. Exp. Neuropsychol.*, 22: 656–676.
- Steinmetz, P.N., Roy, A., Fitzgerald, P.J., Hsiao, S.S., Johnson,
27 K.O. and Niebur, E. (2000) Attention modulates synchro-
nized neuronal firing in primate somatosensory cortex. *Nat-*
29 *ure*, 404: 187–190.
- Sterr, A., Muller, M.M., Elbert, T., Rockstroh, B., Pantev, C.
and Taub, E. (1998) Changed perceptions in Braille readers.
31 *Nature*, 391: 134–135.
- Stryker, M. and Strickland, S. (1984) Physiological segregation
33 of ocular dominance columns depends on the pattern of
afferent electrical activity. *Invest. Ophthalmol. Vis. Sci. Suppl.*,
35 25: 278.
- Sur, M., Garraghty, P.E. and Roe, A.W. (1988) Experimentally
induced visual projections into auditory thalamus and cortex.
37 *Science*, 242: 1437–1441.
- Taub, E. and Uswatte, G. (2003) Constraint-induced movement
therapy: bridging from the primate laboratory to the stroke
39 rehabilitation laboratory. *J. Rehabil. Med.*, 41: 34–40.
- Tegenthoff, M., Cornelius, B., Pleger, B., Malin, J.P. and
Schwenkreis, P. (2004) Amphetamine enhances training-in-
41 duced motor cortex plasticity. *Acta. Neurol. Scand.*, 109:
43 330–336.
- Tobey, E.A., Devous Sr, M.D., Buckley, K., Overson, G., Har-
45 ris, T., Ringe, W. and Martinez-Verhoff, J. (2005) Pharma-
cological enhancement of aural habilitation in adult cochlear
47 implant users. *Ear Hear.*, 26: 45S–56S.

1	Treue, S. and Maunsell, J.H. (1999) Effects of attention on the processing of motion in macaque middle temporal and medial superior temporal visual cortical areas. <i>J. Neurosci.</i> , 19: 7591–7602.	1
3		3
5	Tsodyks, M. (2002) Spike-timing-dependent synaptic plasticity — the long road towards understanding neuronal mechanisms of learning and memory. <i>Trends Neurosci.</i> , 25: 599–600.	5
7		7
9	Walker-Batson, D., Smith, P., Curtis, S. and Unwin, D.H. (2004) Neuromodulation paired with learning dependent practice to enhance post stroke recovery? <i>Restor. Neurol. Neurosci.</i> , 22: 387–392.	9
11	Wallace, H. and Fox, K. (1999) The effect of vibrissa deprivation pattern on the form of plasticity induced in rat barrel cortex. <i>Somatosens. Mot. Res.</i> , 16: 122–138.	11
13	Wang, X., Merzenich, M.M., Sameshima, K. and Jenkins, W.M. (1995) Remodelling of hand representation in adult cortex determined by timing of tactile stimulation. <i>Nature</i> , 378: 71–75.	13
15		15
17		17
19		19
21		21
23		23
25		25
27		27
29		29
31		31
33		33
35		35
37		37
39		39
41		41
43		43
45		45
47		47

Watanabe, T., Nanez, J.E. and Sasaki, Y. (2001) Perceptual learning without perception. *Nature*, 413: 844–848.

Weinberger, N.M. (2003) The nucleus basalis and memory codes: auditory cortical plasticity and the induction of specific, associative behavioral memory. *Neurobiol. Learn Mem.*, 80: 268–284.

Xerri, C., Merzenich, M.M., Peterson, B.E. and Jenkins, W. (1998) Plasticity of primary somatosensory cortex paralleling sensorimotor skill recovery from stroke in adult monkeys. *J. Neurophysiol.*, 79: 2119–2148.

Zhang, Y., Hamilton, S.E., Nathanson, N.M. and Yan, J. (2005). Decreased input-specific plasticity of the auditory cortex in mice lacking M1 muscarinic acetylcholine receptors. *Cereb. Cortex*, In press.

Zhou, Q., Tao, H.W. and Poo, M.M. (2003) Reversal and stabilization of synaptic modifications in a developing visual system. *Science*, 300: 1953–1957.

AUTHOR QUERY FORM

ELSEVIER

Progress in Brain Research, Volume 157

JOURNAL TITLE:	PBR-V157
ARTICLE NO:	57007

Queries and / or remarks

Query No	Details required	Author's response
AQ1	Please provide book title and name of the editors for ref. Carrasco et al. (2004) in the reference list	
AQ2	Please provide the vol. and page range in the ref. Zhang et al. (2005).	
AQ3	The reference McEwen (2003) has not been included in the reference list. Please provide.	
AQ4	In Figure 2, label (a) description is not given. Please include the description for label (a) as for labels (b)-(f).	
AQ5	The reference Kilgard and Merzenich (2001) has not been included in the reference list. Please provide.	
UC REF	If references appear under section "Uncited References", then cite at relevant places in the text. In case of nonavailability of citation, the corresponding references will be deleted from the reference list. Please ignore this query if there is no 'Uncited Reference' section.	

AUTHOR QUERY FORM

ELSEVIER

**Progress in Brain
Research, Volume 157**

JOURNAL TITLE:	PBR-V157
ARTICLE NO:	57007

Queries and / or remarks

Query No	Details required	Author's response