

Selected references

- 1 McGehee, D.S. and Role, L.W. (1995) *Annu. Rev. Physiol.* 57, 521–546
- 2 Colquhoun, L.M. and Patrick, J.W. (1997) *Adv. Pharmacol.* 39, 191–220
- 3 Ramirez-Latorre, J. et al. (1996) *Nature* 380, 347–351
- 4 Wonnacott, S.B. (1997) *Trends Neurosci.* 20, 92–98
- 5 Levin, E.D. and Simon, B.B. (1998) *Psychopharmacology* 138, 217–230
- 6 James, J.R. and Nordberg, A. (1995) *Behav. Genet.* 25, 149–159
- 7 Woolf, N.J. (1991) *Prog. Neurobiol.* 37, 474–524
- 8 Whitehouse, P.J. et al. (1982) *Science* 215, 1237–1239
- 9 Squire, L.R. and Zola-Morgan, S. (1988) *Trends Neurosci.* 11, 170–175
- 10 Eichenbaum, H. et al. (1996) *Proc. Natl. Acad. Sci. U. S. A.* 93, 13500–13507
- 11 Bear, M.F. (1996) *Proc. Natl. Acad. Sci. U. S. A.* 93, 13453–13459
- 12 Frotscher, M. and Léránth, C. (1985) *J. Comp. Neurol.* 239, 237–246
- 13 Matthews, D.A. et al. (1987) *Brain Res.* 402, 30–43
- 14 Jones, S. and Yakel, J.L. (1997) *J. Physiol.* 504, 603–610
- 15 Alkondon, M. et al. (1997) *J. Pharmacol. Exp. Ther.* 283, 1396–1411
- 16 Frazier, C.J. et al. (1998) *J. Neurosci.* 18, 1187–1195
- 17 McQuiston, A.R. and Madison, D.V. (1999) *J. Neurosci.* 19, 2887–2896
- 18 Frazier, C.J. et al. (1998) *J. Neurosci.* 18, 8228–8235
- 19 Gray, R. et al. (1996) *Nature* 383, 713–716
- 20 Jones, S. and Yakel, J.L. (1999) *Cell Biochem. Biophys.* 31, 47–58
- 21 Stewart, M. and Fox, S.E. (1990) *Trends Neurosci.* 13, 163–169
- 22 Fisahn, A. et al. (1998) *Nature* 394, 186–189
- 23 Cobb, S.R. et al. (1999) *J. Physiol.* 518, 131–140
- 24 Yamamoto, J. (1998) *Eur. J. Pharmacol.* 359, 131–133
- 25 Alkondon, M., Pereira, E.F.R. and Albuquerque, E.X. (1998) *Brain Res.* 810, 257–263
- 26 Alkondon, M. et al. (1999) *J. Neurosci.* 19, 2693–2705
- 27 Roerig, B., Nelson, D.A. and Katz, L.C. (1997) *J. Neurosci.* 17, 8353–8362
- 28 Xiang, Z., Huguenard, J.R. and Prince, D.A. (1998) *Science* 281, 985–988
- 29 Sudweeks, S. and Yakel, J.L. (1999) *Soc. Neurosci. Abstr.* 25, 2241
- 30 Picciotto, M.R. et al. (1995) *Nature* 374, 65–67
- 31 Lu, Y. et al. (1998) *J. Pharmacol. Exp. Ther.* 287, 648–657
- 32 Gil, Z., Connors, B.W. and Amitai, Y. (1997) *Neuron* 19, 679–686
- 33 Vidal, C. and Changeaux, J.-P. (1993) *Neuroscience* 56, 23–32
- 34 Bliss, T.V.P. and Collingridge, G.L. (1993) *Nature* 361, 31–39
- 35 Hunter, B.E. et al. (1994) *Neurosci. Lett.* 168, 130–134
- 36 McNaughton, B.L. and Morris, R.G.M. (1987) *Trends Neurosci.* 10, 408–415
- 37 Role, L.W. and Berg, D.K. (1996) *Neuron* 16, 1077–1085
- 38 Zheng, J.Q. et al. (1994) *Nature* 368, 140–144
- 39 Bear, M.F., Carnes, K.M. and Ebner, F.F. (1985) *J. Comp. Neurol.* 237, 519–532
- 40 McDonald, J.K., Speciale, S.G. and Parnavelas, J.G. (1987) *Neuroscience* 21, 825–832
- 41 Henderson, Z. (1991) *Neuroscience* 44, 331–324
- 42 Robertson, R.T. et al. (1991) *Dev. Brain Res.* 58, 81–95
- 43 Kristt, D.A. (1979) *J. Comp. Neurol.* 186, 1–15
- 44 Bear, M.F. and Singer, W. (1986) *Nature* 320, 172–176
- 45 Sato, H. et al. (1987) *J. Neurophysiol.* 58, 765–780
- 46 Gu, Q. and Singer, W. (1993) *Eur. J. Neurosci.* 5, 475–485
- 47 Fuchs, J.L. (1989) *Brain Res.* 501, 223–234
- 48 Broide, R.S. et al. (1995) *Neuroscience* 67, 83–94
- 49 Aramakis, V.B. and Metherate, R. (1998) *J. Neurosci.* 18, 8485–8495
- 50 Robinson, T.E. and Berridge, K.C. (1993) *Brain Res. Rev.* 18, 247–291
- 51 Kalivas, P.W. and Stewart, J. (1991) *Brain Res. Rev.* 16, 223–244
- 52 Stolerman, I.P. and Shoab, M. (1991) *Trends Pharmacol. Sci.* 12, 467–473
- 53 Corrigan, W.A. (1991) *Br. J. Addiction* 86, 507–510
- 54 Corrigan, W.A. et al. (1992) *Psychopharmacology* 107, 285–289
- 55 Nisell, M., Nomikos, G.G. and Svensson, T.H. (1995) *Pharmacol. Toxicol.* 76, 157–162
- 56 Nisell, M., Nomikos, G.G. and Svensson, T.H. (1994) *Pharmacol. Toxicol.* 75, 348–352
- 57 Nisell, M., Nomikos, G.G. and Svensson, T.H. (1994) *Synapse* 16, 36–44
- 58 Yeomans, J.S. (1995) *Neuropsychopharmacology* 12, 3–16
- 59 Blaha, C.D. (1996) *J. Neurosci.* 16, 714–722
- 60 Futami, T., Takakusaki, K. and Kitai, S.T. (1995) *Neurosci. Res.* 21, 331–342
- 61 Grenhoff, J., Aston-Jones, G. and Svensson, T.H. (1986) *Acta. Physiol. Scand.* 128, 351–358
- 62 Wonnacott, S. (1990) *Trends Pharmacol. Sci.* 11, 216–219
- 63 Dani, J.A. and Heinemann, S. (1996) *Neuron* 16, 905–908
- 64 Calabresi, P., Lacey, M.G. and North, R.A. (1989) *Br. J. Pharmacol.* 98, 135–140
- 65 Pidoplichko, V.I. et al. (1997) *Nature* 390, 401–404
- 66 Picciotto, M.R. et al. (1998) *Nature* 391, 173–177
- 67 Shoab, M. et al. (1994) *Br. J. Pharmacol.* 111, 1073–1080
- 68 Schilström, B. et al. (1998) *Neuroscience* 82, 781–789
- 69 Wolf, M.E. (1998) *Prog. Neurobiol.* 54, 679–720

Acknowledgements

The authors thank D.L. Armstrong, D. Pettit and S.D. Bryant for their insightful comments on this manuscript.

Long-term potentiation in the amygdala: a mechanism for emotional learning and memory

Stephen Maren

In the mammalian brain, LTP is an enduring form of synaptic plasticity that is posited to have a role in learning and memory. Compelling new evidence for this view derives from studies of LTP in the amygdala, a brain structure that is essential for simple forms of emotional learning and memory, such as Pavlovian fear conditioning in rats. More specifically, antagonists of the NMDA receptor block both amygdaloid LTP induction and fear conditioning, fear conditioning induces increases in amygdaloid synaptic transmission that resemble LTP, and genetic modifications that disrupt amygdaloid LTP eliminate fear conditioning. Collectively, these results provide the most-convincing evidence to date that LTP mediates learning and memory in mammals.

Trends Neurosci. (1999) 22, 561–567

THE CAPACITY to learn that some events in the world have adverse consequences is not only practical, but is also adaptive: individuals that cannot learn to recognize threats in the environment are doomed to injury

and death. Pavlovian fear conditioning is a simple form of associative learning that supports the acquisition of this type of information. Simply stated, fear conditioning occurs when an innocuous stimulus (the conditional

Stephen Maren is at the Dept of Psychology and Neuroscience Program, University of Michigan, MI 48109-1109, USA.




Fig. 1. Anatomy and connectivity of fear-conditioning circuits in the rat amygdala. As shown in the thionin-stained coronal section (A), and enlarged in the inset, the amygdala is a complex of nuclei located deep within the temporal lobe. The colored overlays highlight the amygdaloid nuclei known to be important in fear conditioning. These include the lateral (LA), basolateral (BL) and basomedial (BM) nuclei, which together form the basolateral complex (BLA), and the central nucleus, CEA (CEM, medial and CEI, lateral). The BLA receives and integrates sensory information from a variety of sources [arrows in (B)], including the thalamic medial geniculate nucleus (MGN, auditory), the perirhinal cortex (PERI, visual), the insular cortex (INS, gustatory and somatosensory), the thalamic posterior intralaminar nucleus (PIL, somatosensory), the hippocampal formation (HIP, spatial and contextual) and piriform cortex (PIR, olfactory). Thus, the BLA is a locus of sensory convergence and a site for conditional stimulus–unconditional stimulus (CS–US) association within the amygdala. Intra-amygdaloid circuitry conveys the CS–US association to the CEA, where divergent projections to the hypothalamus and brainstem mediate fear responses such as freezing (periaqueductal gray, PAG), potentiated acoustic startle (nucleus reticularis pontis caudalis, RPC), increased heart rate and blood pressure (lateral hypothalamus, LH), increased respiration (parabrachial nucleus, PB) and glucocorticoid release (paraventricular nucleus of the hypothalamus, PVN). Scale bars, 1 mm.

trends in Neurosciences

stimulus or CS) precedes the occurrence of a different, aversive stimulus (the unconditional stimulus or US). In rats, pairing a neutral CS, such as a loud tone, with an aversive US, such as an electric footshock, generates fear conditioning. After a few trials, the CS eventually elicits robust conditional fear responses (CRs), which include potentiated acoustic startle, hypoalgesia, elevated blood pressure and freezing (immobility except for breathing). Fear conditioning is acquired rapidly, retained over long periods of time, and is readily studied in both rats and humans. Indeed, the ease of establishing and assessing fear memories has established fear conditioning as a popular learning task for behavioral, biochemical and molecular studies of learning and memory.

Because of the relevance of fear conditioning to disorders of fear and anxiety in humans, a major goal of research has been to identify the crucial brain substrates and neuronal mechanisms that underlie Pavlovian fear conditioning in rats. After years of research, there is almost unanimous agreement that the amygdala, a nut-shaped structure that lies deep within the temporal lobes, is essential for forming and storing fear memories^{1–4}. The identification of an essential basis for fear conditioning has now opened the door to studies of synaptic and molecular processes within the amygdala that underlie this form of learning. One candidate cellular mechanism for amygdala-dependent learning is LTP, an enduring form of synaptic plasticity that was discovered in the hippocampus and has recently been identified in the amygdala. In this article, the compelling new evidence that implicates amygdaloid LTP in fear conditioning will be described, and the possibility that synaptic

potentiation in the amygdala is a mechanism for emotional learning and memory will be discussed. The focus on fear conditioning in this article should not overshadow the other roles posited for the amygdala, such as the modulation of declarative memory consolidation⁵ and attention⁶.

Neural circuitry of fear conditioning

It has been recognized for decades that the amygdala is involved in emotional processes⁷, including aversively motivated learning^{8,9} (see Ref. 10 for an excellent review of amygdaloid anatomy). The major amygdaloid nuclei and projections are shown in Fig. 1. Recent anatomical and behavioral evidence indicates that there are at least two distinct subsystems within the amygdala that are important for Pavlovian fear conditioning^{11,12}. The first subsystem of the amygdala consists of the lateral, basolateral and basomedial nuclei. These nuclei, which are collectively referred to as the basolateral amygdaloid complex (BLA), form the primary sensory interface of the amygdala. Thus, the BLA receives synaptic input from many primary sensory structures, and lesions in

these structures yield deficits in Pavlovian fear conditioning. For example, projections from the auditory thalamus and auditory cortex to the BLA are essential for conditioning to auditory CSs (Refs 13–15), projections from the hippocampal formation to the BLA appear to underlie conditioning to contextual CSs (Refs 16–18), and projections from the perirhinal cortex transmit visual CS information to the BLA (Ref. 19). Information about the aversive footshock US might reach the BLA via parallel thalamic and cortical pathways²⁰. Consistent with this anatomy, single neurons in the BLA respond to auditory, visual and somatic (shock) stimuli, which indicates that the amygdala is a locus of convergence for information about CSs and USs (Ref. 21). The BLA is, therefore, anatomically situated to integrate information from a variety of sensory domains.

The information processed by the BLA is either relayed back to afferent structures or sent to the second major subsystem of the amygdala, the central nucleus of the amygdala (CEA). The CEA projects to many brainstem targets and is the amygdala's interface with the fear-response systems. For example, the CEA projects to the nuclei in the hypothalamus, midbrain and medulla that control a variety of defensive responses, including freezing and acoustic startle. Electrical stimulation of the CEA produces responses that are similar to those elicited by stimuli paired with shock^{22,23}, and lesions of the CEA also produce profound deficits in both the acquisition and expression of conditional fear^{24,25}. Moreover, lesions placed in structures that are efferent to the CEA, such as the lateral hypothalamus or periaqueductal grey (PAG), produce selective deficits in either cardiovascular or somatic conditional fear responses, respectively²⁶. This

suggests that the CEA is the final common pathway for the generation of learned fear responses. Thus, the amygdala contains two distinct subsystems that represent areas of either sensory convergence (BLA) or response divergence (CEA).

Fear memories are formed and stored in the basolateral amygdala

Much evidence indicates that the BLA is the crucial neural locus for the formation and storage of fear memories. Selective lesions of the BLA abolish both acquisition and expression of conditional fear in several behavioral paradigms^{27,28}. In addition, BLA lesions yield deficits in conditional fear when they are made up to one month after training^{29,30} or after extensive overtraining³¹. Similar to permanent lesions, manipulations that temporarily disable amygdaloid neurons prevent both the acquisition and expression of fear conditioning^{32,33}. Fear-conditioning deficits are associative in nature, because rats with BLA lesions can perform the freezing response under some conditions³⁴. Consistent with this, reversible inactivation of the BLA during training produces learning deficits that are manifest during testing with an intact amygdala^{32,33}.

The view that the BLA is a locus of plasticity during aversive learning is further supported by electrophysiological studies of neuronal activity in the amygdala during auditory fear conditioning. For example, neurons in the amygdala exhibit short-latency, CS-elicited firing during aversive learning^{35–37}. Associative neuronal firing in the BLA precedes the development of both behavioral CRs and associative firing in other brain structures, including the auditory cortex³⁸. Moreover, an intact amygdala is required for the acquisition of at least some forms of neuronal plasticity in the auditory cortex³⁹. Because of its essential role in forming and storing fear memories, the BLA serves as an ideal anatomical substrate for analyzing the relationship of synaptic plasticity mechanisms, such as LTP, to behavior^{12,40}.

Amygdaloid neurons exhibit LTP

The first examination of synaptic plasticity in the amygdala was performed by Racine and colleagues in awake, behaving rats. In this preparation, 'tetanic' LTP (which is contrasted with 'behavioral' LTP) was induced in the amygdala by applying high-frequency electrical stimulation to the pyriform cortex⁴¹. This LTP consisted of a long-term enhancement in the amplitude of stimulus-elicited extracellular field potentials recorded in the amygdala. More recently, tetanic LTP has been reported in both thalamo-amygdala and hippocampo-amygdala projections *in vivo*^{42–44}. Fig. 2 illustrates the latter. Both of these pathways are believed to transmit CS information to BLA neurons, and potentiation of thalamo-amygdaloid synapses enhances the amplitude of auditory-stimulus-elicited potentials in the BLA




Fig. 2. Tetanic LTP is induced in the basolateral nucleus of the amygdala by electrical stimulation of the hippocampal formation. Stimulating electrodes were placed in the ventral hippocampus (light gray) and a recording electrode was placed in the basolateral amygdala (dark gray) of anesthetized rats (A). (B) Single-pulse stimulation of the ventral hippocampus elicited a biphasic extracellular field potential (traces) with a short-latency negative-going component (indicated by the black dots). The potentials obtained following high-frequency and low-frequency stimulation are shown on the left and right, respectively. (C) High-frequency stimulation (black), but not low-frequency stimulation (gray), of the hippocampal formation (at the times indicated by the arrows in the line plot) induced a robust LTP of the short-latency component of the amygdaloid field potential (pre- and post-tetanus responses are superimposed). Adapted, with permission, from Ref. 42.

(Ref. 44). Studies using intracellular recording in amygdala slices have confirmed that BLA neurons exhibit tetanic LTP *in vitro*⁴⁵.

As in the hippocampus^{46,47}, LTP induction in the BLA has been reported to require the activation of NMDA receptors^{42,48,49}. However, there are indications that LTP induction in the amygdala does not require NMDA-receptor activation under some conditions^{48,50}. In fact, LTP induction on amygdaloid interneurons appears to be mediated by Ca²⁺-permeable AMPA receptors, rather than NMDA receptors⁵¹. This suggests that there are both NMDA-receptor-dependent and NMDA-receptor-independent forms of LTP in the amygdala. Once induced, the mechanisms involved in expressing amygdaloid LTP are not fully understood. There is evidence for increased presynaptic neurotransmitter release following LTP induction in the amygdala (that is, paired-pulse facilitation in the BLA is attenuated following LTP induction)^{42,48}, although changes in the properties of postsynaptic AMPA receptors cannot be ruled out⁴². It should also be noted that BLA neurons also exhibit LTD of synaptic transmission under some conditions and metabotropic glutamate receptors appear to regulate the induction of this form of plasticity⁵². Although not yet tested, LTD at synapses in the amygdala might also be involved in fear conditioning, particularly with regard to the acquisition of inhibitory associations.

Glutamate receptors in the amygdala are essential for fear conditioning

The important role for glutamate receptors in amygdaloid LTP suggests a role for these receptors in the acquisition and expression of Pavlovian fear conditioning. Consistent with this view, several laboratories have now demonstrated that the infusion of NMDA-receptor antagonists, such as (\pm)-2-amino-5-phosphonovaleric




Fig. 3. Behavioral LTP is induced in the lateral nucleus of the amygdala by Pavlovian fear conditioning. A recording electrode was placed in the lateral nucleus of the amygdala (LA, dark gray) to measure auditory-stimulus-elicited extracellular field potentials in awake rats (A). (B) Brief tone stimuli elicited a biphasic extracellular field potential with a short-latency negative component (pre-training, indicated by the black dots). (C) Pairing the tone stimulus with an aversive footshock (training) potentiated the short-latency negative component, and this potentiation persisted during a shock-free extinction test (testing). Potentiation of the LA field potential did not occur in rats receiving unpaired tones and footshocks. (D) Both paired (black) and unpaired (gray) training generated freezing during the conditioning phase, although paired training induced significantly greater freezing to the tone CS than unpaired training during testing. Adapted, with permission, from Ref. 62.

acid (APV), into the BLA prevents the acquisition of conditional fear responses, including freezing⁵³⁻⁵⁵ and fear-potentiated startle^{56,57}. Moreover, it has been reported that AMPA-receptor antagonists impair the expression of conditional fear⁵⁸ and that AMPA-receptor agonists enhance the acquisition of fear conditioning⁵⁹. These results reveal an important role for amygdaloid glutamate receptors in the acquisition and expression of fear conditioning.

One unanticipated finding that has emerged from this work is that pre-testing infusions of APV into the amygdala impair the expression of some learned fear responses, such as freezing^{54,55}, but leaves others, such as fear-potentiated startle, intact^{56,57}. The disruption of conditional freezing might be explained, at least in part, by the observation that APV reduces cell excitability at some synapses in the amygdala *in vivo*^{42,60}. Moreover, it is possible that conditional fear responses, such as freezing and fear-potentiated startle, are under the control of two different populations of amygdaloid neurons that differ in their levels of NMDA receptors⁵⁴. Double-labeling studies that have examined NMDA-receptor binding in neurons in the amygdala that project to the PAG and nucleus reticularis pontis caudalis (RPC), which control freezing and startle responses, respectively, would be useful for investigating this possibility. Even so, the similar effects of APV on both the acquisition and expression of conditional freezing is problem-

atic for linking amygdaloid NMDA receptors to fear conditioning, insofar as NMDA receptors are only thought to have a role in encoding, not retrieving, fear memories. One might be tempted to conclude that NMDA-receptor antagonists exert their action on fear conditioning through a reduction in amygdaloid cell excitability, rather than a disruption of LTP induction. While this might be true for conditional freezing, the selective effect of intra-amygdala administration of APV on the acquisition of fear-potentiated startle, as opposed to the expression, suggests that APV retards fear conditioning through an attenuation of amygdaloid LTP rather than a reduction in cell excitability⁶¹.

Fear conditioning induces LTP in the amygdala

The ability of NMDA-receptor antagonists to block both the induction of tetanic LTP in the amygdala and the acquisition of fear conditioning suggests that a process akin to tetanic LTP might underlie the acquisition of conditional fear in awake, behaving rats. Two recent studies are consistent with this possibility. In the first study, LeDoux and colleagues found that fear conditioning in response to an auditory CS enhanced auditory-elicited potentials in the BLA in a manner that was similar to that observed following the induction of tetanic LTP in

auditory projections to the BLA (see Fig. 3)⁶². In other words, both behavioral learning and high-frequency electrical stimulation enhanced the amplitude of extracellular auditory-stimulus-elicited field potentials in the amygdala. The similar potentiation of auditory-stimulus-elicited field potentials by high-frequency stimulation and behavioral learning suggests that LTP in auditory pathways to the amygdala might in fact mediate fear conditioning to an auditory CS.

The second study that has examined the relationship between behavioral learning and amygdaloid LTP used a different approach. McKernan and colleagues first trained rats in a fear-potentiated startle paradigm and then sacrificed the animals in order to obtain intracellular recordings from amygdaloid neurons in brain slices⁶³. One advantage of this preparation is that the investigators could examine pharmacologically characterized synaptic currents in single amygdaloid neurons. These investigators found that prior fear conditioning selectively enhanced excitatory AMPA-receptor-mediated synaptic currents in thalamic projections to BLA neurons, but not in projections from the endopyriform nucleus. AMPA-receptor-mediated currents were not enhanced in rats that received unpaired tones and footshocks (a control procedure for the non-associative effects of tone and footshock). NMDA-receptor-mediated currents were not potentiated under any circumstances. These data indicate that associative

learning in a fear-conditioning paradigm can produce a selective augmentation of excitatory synaptic transmission in auditory projections to the amygdala.

McKernan and colleagues provided some clues to the nature of the synaptic modification that underlies the enhanced AMPA-receptor-mediated currents in BLA neurons with an analysis of paired-pulse facilitation of excitatory synaptic currents recorded from amygdaloid neurons⁶³. As has been reported following the induction of tetanic LTP in the amygdala^{42,48}, rats that underwent paired training exhibited reduced paired-pulse facilitation⁶³. This suggests that both behavioral LTP and tetanic LTP are associated with increases in neurotransmitter release, insofar as manipulations that increase neurotransmitter release (such as decreasing the extracellular $[Mg^{2+}]/[Ca^{2+}]$ ratio) also reduce paired-pulse facilitation. Paired-pulse facilitation is not attenuated in endopyriform projections to BLA neurons.

Taken together, this pair of studies^{62,63} indicates that fear conditioning induces behavioral LTP in the amygdala. This outcome was apparent whether synaptic responses were measured 'on-line' in the awake, behaving rat⁶² or 'off-line' in amygdala slices obtained from previously conditioned rats⁶³. In both preparations, behavioral LTP exhibited properties that were common to tetanic LTP. For example, both behavioral and tetanic LTP were manifest in enhanced auditory-stimulus-elicited potentials *in vivo*, and reduced paired-pulse facilitation *in vitro*. Although it has not yet been examined, an important issue concerns the NMDA-receptor dependency of behavioral LTP in these preparations. A similar dependency of both behavioral LTP induction and fear conditioning on NMDA-receptor activation would provide further support for the involvement of amygdaloid LTP in the acquisition of fear memories.

Molecular basis of amygdaloid LTP and fear conditioning

A novel and exciting approach to examining the relationship between synaptic plasticity mechanisms, and learning and memory involves the use of genetically modified animals. In the past few years, many mice have been engineered with genetic manipulations that either disable or eliminate key proteins in the intracellular biochemical cascades that mediate LTP. While these studies have largely focused on the relationship of hippocampal LTP to behavior, several recent investigations using genetically modified mice have probed the link between amygdaloid LTP and Pavlovian fear conditioning.

Indirect evidence for a role for amygdaloid LTP in fear conditioning has come from the work of Kandel and colleagues who used mice that overexpressed a mutant




Fig. 4. Mice that lack RasGRF exhibit deficits in both tetanic LTP in the amygdala and Pavlovian fear conditioning. Tetanic LTP in the basolateral nucleus of the amygdala (BL) was characterized *in vitro* in brain slices obtained from wild-type mice and mice lacking RasGRF. Extracellular field potentials in BL were elicited by electrical stimulation of the lateral nucleus of the amygdala (LA) (A). High-frequency stimulation of LA induced a robust LTP of synaptic transmission in the BL of wild-type mice (black), but resulted in a rapidly decaying potentiation in mice that lack RasGRF (gray) (B). Consistent with the deficit in tetanic LTP in the amygdala, mice lacking RasGRF exhibited deficits in long-term retention of Pavlovian fear conditioning (C). Conditional freezing to a tone that was paired with footshock (left) and the context in which the tone-shock pairing occurred (right) was impaired in RasGRF knockouts when they were tested 24 h after training. However, their short-term memory for the fear conditioning was intact when they were tested 30 min after training. This indicates that the deficit in long-term retention in mice lacking RasGRF was not the result of an inability of these mice to exhibit freezing behavior. Hippocampal LTP in RasGRF knockouts was normal (data not shown). Scale bar, 1 mm. Adapted, with permission, from Ref. 69.

form of the gene for Ca²⁺/calmodulin-dependent protein kinase II (CaMKII) in the amygdala, hippocampus and striatum⁶⁴. In the hippocampus, CaMKII is activated by Ca²⁺ influx through the NMDA receptor. This kinase interacts with a number of substrates including the cAMP-response-element-binding protein (CREB), a nuclear transcription factor that has been implicated in learning and memory. Not surprisingly, CaMKII has been demonstrated to have a role in LTP induction, and mice that overexpress the transgene for this protein do not exhibit LTP in the hippocampus following theta-frequency stimulation, for example⁶⁵. Interestingly, these transgenic mice, as well as mice that lack CaMKII, also exhibit impairments in Pavlovian fear conditioning to both contextual and auditory cues^{64,66}. A similar pattern has been observed in mice with a targeted disruption of CREB, which has been implicated in hippocampal LTP (Ref. 67), and whose synthesis is induced in both the amygdala and hippocampus by fear conditioning⁶⁸. The global impairment in conditional fear to both contextual and auditory cues suggests the existence of amygdala dysfunction in these genetically modified mice. Although the nature of this dysfunction is not yet known, it is tempting to speculate that it will take the form of impaired amygdaloid LTP.

A more-direct demonstration of the role for amygdaloid LTP in fear conditioning comes from studies of mice that lack RasGRF, a neuron-specific guanine nucleotide releasing factor that is activated by both Ca²⁺ and G-protein-coupled messengers⁶⁹. As shown in Fig. 4, electrophysiological recordings from brain slices

obtained from such mice reveal a pronounced deficit in LTP in the basolateral nucleus of the amygdala. Interestingly, these mice also exhibit impairments in consolidating long-term memories for Pavlovian fear conditioning to both contextual and acoustic stimuli (see Fig. 4). These deficits in LTP and learning were selective for the amygdala and Pavlovian fear conditioning insofar as both hippocampal LTP and spatial learning in RasGRF knockouts were normal. Interestingly, RasGRF modulates CREB activity through the mitogen-activated protein kinase (MAPK) pathway. A role for MAPK in fear conditioning has recently been demonstrated⁷⁰. Together, these results provide strong support for the view that synaptic LTP in the amygdala is required for the establishment and maintenance of emotional memories.

Associative LTP as a mechanism for Pavlovian fear conditioning

If LTP in the amygdala is a synaptic mechanism for the acquisition of conditional fear, then it is of interest to consider how such a mechanism might operate during learning. In the hippocampus, tetanic LTP exhibits a property known as associativity (for example, see Ref. 71). Thus, LTP can be induced in 'weak' synaptic pathways if activity in these pathways is paired with activity in a 'strong' pathway. This associative property of LTP can accommodate CS-US association formation in the amygdala during Pavlovian fear conditioning^{4,40,72}. For example, in the conditional freezing paradigm, an auditory cue does not strongly activate amygdaloid neurons³⁸ and, therefore, does not elicit freezing prior to fear conditioning. In contrast, we assume that the footshock US strongly activates amygdaloid neurons to produce a central fear state. Paradoxically, activation of the amygdala by shock USs does not translate into a freezing response. This is believed to be true because the footshock US also activates the dorsal PAG, which results in both the shock-elicited activity burst and inhibition of the ventral PAG, an essential structure for generating the freezing response. During fear conditioning, however, activity in the 'weak' CS pathway is paired with depolarization generated by the 'strong' US pathway. Thus, it is through the temporal association of the CS and US that synapses in the CS pathway are potentiated, presumably through a mechanism akin to associative LTP. Although this mechanism appears plausible, further studies are required to determine whether amygdaloid neurons exhibit associative LTP.

Of course, such a simple model cannot account for the richness of learning phenomena in Pavlovian paradigms by itself. Indeed, several investigators have raised doubts about the heuristic value of using LTP as a model for learning and memory⁷³⁻⁷⁵. One argument that has been proposed is that the temporal requirements for associative LTP induction are not congruent with optimal Pavlovian-conditioning parameters. However, in cases where there is divergence in the optimal-stimulus timing parameters for LTP induction and fear conditioning, such as in long-delay fear conditioning, one can certainly imagine mechanisms in the brain that reunite temporally discontinuous events under conditions that will favor associative LTP. In fact, Brown and colleagues⁷⁶ recently proposed a computational model that bridges the long CS-US delays that are typically used in fear-conditioning experiments. Their model is based on the discovery of late-firing neurons⁷⁷ (neurons that withhold

firing an action potential for up to several seconds following injection of a depolarizing current) in the perirhinal cortex, a putative CS pathway to the amygdala. In the model, these late-firing neurons maintain CS-elicited firing during the CS-US delay interval so that the CS and US can be instantiated as contiguous events by amygdala neurons using standard Hebbian modification rules. Thus, associative LTP in the amygdala remains a plausible mechanism for fear conditioning, despite the parametric differences between conditioning parameters and optimal LTP induction parameters. This work reveals that an understanding of the computational properties of amygdaloid circuits provides an important insight into the role synaptic plasticity has in the acquisition of emotional learning and memory.

Concluding remarks

An abundance of evidence indicates that neurons in the amygdala are essential for simple forms of emotional learning and memory, such as Pavlovian fear conditioning in rats. In the amygdala, LTP is an enduring form of synaptic plasticity that has been posited to have a role in Pavlovian fear conditioning. As this article illustrates, this view is supported by compelling new evidence that NMDA-receptor antagonists block both amygdaloid LTP induction and fear conditioning, fear conditioning induces increases in amygdaloid synaptic transmission that resemble LTP, and genetic modifications that eliminate amygdaloid LTP also impair fear conditioning. Collectively, these results provide the strongest evidence to date that LTP mediates learning and memory in mammals.

Selected references

- 1 LeDoux, J. (1998) *Biol. Psychiatry* 44, 1229-1238
- 2 Davis, M. (1997) *J. Neuropsychiatry Clin. Neurosci.* 9, 382-402
- 3 Fanselow, M.S. (1994) *Psychonomic Bull. Rev.* 1, 429-438
- 4 Maren, S. (1996) *Mol. Neurobiol.* 13, 1-22
- 5 McGaugh, J.L. *et al.* (1996) *Proc. Natl. Acad. Sci. U. S. A.* 93, 13508-13514
- 6 Gallagher, M. and Holland, P.C. (1994) *Proc. Natl. Acad. Sci. U. S. A.* 91, 11771-11776
- 7 Klüver, H. and Bucy, P.C. (1937) *Am. J. Physiol.* 119, 352-353
- 8 Brady, J.V. *et al.* (1954) *J. Comp. Physiol. Psychol.* 47, 179-186
- 9 Kelliecutt, M.H. and Schwartzbaum, J.S. (1963) *Psychol. Rep.* 12, 351-358
- 10 Swanson, L.W. and Petrovich, G.D. (1998) *Trends Neurosci.* 21, 323-331
- 11 LeDoux, J.E. (1995) *Annu. Rev. Psych.* 46, 209-235
- 12 Davis, M. *et al.* (1994) *Trends Neurosci.* 17, 208-214
- 13 Campeau, S. and Davis, M. (1995) *J. Neurosci.* 15, 2312-2327
- 14 LeDoux, J.E. *et al.* (1990) *J. Neurosci.* 10, 1062-1069
- 15 Romanski, L.M. and LeDoux, J.E. (1992) *J. Neurosci.* 12, 4501-4509
- 16 Kim, J.J. and Fanselow, M.S. (1992) *Science* 256, 675-677
- 17 Maren, S. *et al.* (1997) *Behav. Brain Res.* 88, 261-274
- 18 Phillips, R.G. and LeDoux, J.E. (1992) *Behav. Neurosci.* 106, 274-285
- 19 Rosen, J.B. *et al.* (1992) *J. Neurosci.* 12, 4624-4633
- 20 Shi, C. and Davis, M. (1999) *J. Neurosci.* 19, 420-430
- 21 Romanski, L.M. *et al.* (1993) *Behav. Neurosci.* 107, 444-450
- 22 Iwata, J. *et al.* (1987) *Brain Res.* 418, 183-188
- 23 Kapp, B.S. *et al.* (1982) *Brain Res.* 234, 251-262
- 24 Hitchcock, J. and Davis, M. (1986) *Behav. Neurosci.* 100, 11-22
- 25 Kim, M. and Davis, M. (1993) *Behav. Neurosci.* 107, 1088-1092
- 26 LeDoux, J.E. *et al.* (1988) *J. Neurosci.* 8, 2517-2529
- 27 Campeau, S. and Davis, M. (1995) *J. Neurosci.* 15, 2301-2311
- 28 Sananes, C.B. and Davis, M. (1992) *Behav. Neurosci.* 106, 72-80
- 29 Maren, S. *et al.* (1996) *Behav. Neurosci.* 110, 718-726
- 30 Lee, Y. *et al.* (1996) *Behav. Neurosci.* 110, 836-839
- 31 Maren, S. (1998) *J. Neurosci.* 18, 3088-3097
- 32 Muller, J. *et al.* (1997) *Behav. Neurosci.* 111, 683-691
- 33 Helmstetter, F.J. and Bellgowan, P.S. (1994) *Behav. Neurosci.* 108, 1005-1009
- 34 Maren, S. (1999) *J. Neurosci.* 19, 8696-8703
- 35 Quirk, G.J. *et al.* (1995) *Neuron* 15, 1029-1039
- 36 Maren, S. *et al.* (1991) *Brain Res.* 549, 311-316

- 37 Pascoe, J.P. and Kapp, B.S. (1985) *Behav. Brain Res.* 16, 117–133
 38 Quirk, G.J. *et al.* (1997) *Neuron* 19, 613–624
 39 Armony, J.L. *et al.* (1998) *J. Neurosci.* 18, 2592–2601
 40 Maren, S. and Fanselow, M.S. (1996) *Neuron* 16, 237–240
 41 Racine, R.J. *et al.* (1983) *Brain Res.* 260, 217–231
 42 Maren, S. and Fanselow, M.S. (1995) *J. Neurosci.* 15, 7548–7564
 43 Clugnet, M.C. and LeDoux, J.E. (1990) *J. Neurosci.* 10, 2818–2824
 44 Rogan, M.T. and LeDoux, J.E. (1995) *Neuron* 15, 127–136
 45 Chapman, P.F. *et al.* (1990) *Synapse* 6, 271–278
 46 Bliss, T.V.P. and Collingridge, G.L. (1993) *Nature* 361, 31–39
 47 Maren, S. and Baudry, M. (1995) *Neurobiol. Learn. Mem.* 63, 1–18
 48 Huang, Y.Y. and Kandel, E.R. (1998) *Neuron* 21, 169–178
 49 Gean, P.W. *et al.* (1993) *Brain Res. Bull.* 31, 7–11
 50 Chapman, P.F. and Bellavance, L.L. (1992) *Synapse* 11, 310–318
 51 Mahanty, N.K. and Sah, P. (1998) *Nature* 394, 683–687
 52 Li, H. *et al.* (1998) *J. Neurosci.* 18, 1662–1670
 53 Fanselow, M.S. *et al.* (1994) *Behav. Neurosci.* 108, 235–240
 54 Lee, H. and Kim, J.J. (1998) *J. Neurosci.* 18, 8444–8454
 55 Maren, S. *et al.* (1996) *Behav. Neurosci.* 110, 1365–1374
 56 Campeau, S. *et al.* (1992) *Behav. Neurosci.* 106, 569–574
 57 Miserendino, M.J. *et al.* (1990) *Nature* 345, 716–718
 58 Kim, M. *et al.* (1993) *Behav. Neural Biol.* 59, 5–8
 59 Rogan, M.T. *et al.* (1997) *J. Neurosci.* 17, 5928–5935
 60 Li, X.F. *et al.* (1995) *Exp. Brain Res.* 105, 87–100
 61 Gewirtz, J.C. and Davis, M. (1997) *Nature* 388, 471–474
 62 Rogan, M.T. *et al.* (1997) *Nature* 390, 604–607
 63 McKernan, M.G. and Shinnick-Gallagher, P. (1997) *Nature* 390, 607–611
 64 Mayford, M. *et al.* (1996) *Science* 274, 1678–1683
 65 Bach, M.E. *et al.* (1995) *Cell* 81, 905–915
 66 Chen, C. *et al.* (1994) *Science* 266, 291–294
 67 Bourchouladze, R. *et al.* (1994) *Cell* 79, 59–68
 68 Impey, S. *et al.* (1998) *Nat. Neurosci.* 1, 595–601
 69 Brambilla, R. *et al.* (1997) *Nature* 390, 281–286
 70 Atkins, C.M. *et al.* (1998) *Nat. Neurosci.* 1, 602–608
 71 Barrionuevo, G. and Brown, T.H. (1983) *Proc. Natl. Acad. Sci. U. S. A.* 80, 7347–7351
 72 Fanselow, M.S. (1993) *Curr. Dir. Psychol. Sci.* 2, 152–156
 73 Diamond, D.M. and Rose, G.M. (1994) *Psychobiology* 22, 263–269
 74 Gallistel, C.R. (1995) in *Brain and Memory: Modulation and Mediation of Neuroplasticity* (McGaugh, J.L., Weinberger, N.M. and Lynch, G., eds), pp. 328–337, Oxford University Press
 75 Shors, T.J. and Matzel, L.D. (1997) *Behav. Brain Sci.* 20, 597–634
 76 Tieu, K.H. *et al.* (1999) *Psychobiology* 27, 1–25
 77 Faulkner, B. and Brown, T.H. (1999) *J. Comp. Neurol.* 411, 613–642

Acknowledgements
 The author's research was supported by a grant from the National Institute of Mental Health (R29MH57865). The author thanks two anonymous referees for helpful comments on a previous version of this manuscript.

BOOK REVIEWS

States of Mind: New Discoveries about How Our Brains Make Us Who We Are

edited by Roberta Conlan, John Wiley & Sons, 1999. £19.95 (x + 214 pages)
 ISBN 0 471 29963 4

Dissemination is the soul of science. Not only is it even more vital that we inform an increasingly sceptical public about what we are doing, but a sufficient number of teenagers need to be excited by it. I am almost sure that reading the right article at the right time is a life event that seeds some future crucial bias to study the brain rather than rocket science. So any attempt to fire such enthusiasm by individual scientists sometimes, as here in collaboration with educational organizations such as the Dana Alliance, deserves our fullest admiration.

What is the best way to disseminate? The televised Christmas Royal Institution Lectures in the UK have enormous and well-deserved impact. *States of Mind* arises from a series of lectures given by distinguished neuroscientists in 1997, in the USA. Whether a 'book of the lectures' is actually the best way to disseminate the material to a wider audience is debatable. It represents a tall order for the erstwhile contributors: it is one thing to provide an entertaining and simplified lecture using good visuals, but quite another to consolidate it to the written page. The assiduous copy editor will badger you into clarifying that minor ambiguity into which you are tempted merely to protect yourself from accusations of opting for the simple story. Your scientific colleagues are liable to read the material in a perversely critical way, especially if you have not conveyed their important theory with complete precision or, worse still, you have omitted to cite them altogether. For an editor, the task of knowing which themes to pursue in such a book, and how optimally to harmonize

and sequence the varied contributions from several authors, is a major challenge.

States of Mind rises some way to these challenges. The book focuses on the neural and neurochemical basis of behaviour and individual differences. The title is slightly misleading as the major state of mind, viz consciousness, is alluded to only briefly. There is a stimulating exposition of arousal states in a chapter on sleep by Hobson. The distinction between implicit and explicit memory invokes consciousness in two of the chapters. But most of the advances described are most relevant to the unconscious state. Kandel takes a clearly argued reductionist line on the molecular basis of learning by reference to his famous work on *Aplysia*. However, for novices it must be disconcerting that this relates to the 'wastebasket' category of forms of learning (termed 'non-declarative memory') rather than the exciting 'declarative' memory deficits they have just learned about from the earlier account of the patient, H.M. Thus, the vaunted synthesis of molecular neuroscience with cognitive psychology is not necessarily portrayed in perhaps the most advantageous light. The implications of the reductionist approach to memory seem very clear; it will indeed be exciting if the same basic molecular mechanisms operate in declarative as well as non-declarative memory. However, we will then have to understand the undoubted differences in functioning of such memory systems in other ways; for example at the computational or neural systems levels, if we are to appreciate their adaptive significance. What does come over well in this context

is LeDoux's cameo of his excellent popular work¹: the possible interplay between implicit memory and working-memory systems in emotion is discussed lucidly, for example, alongside the use of 'neuroscience tools' to investigate the amygdala.

However, it would perhaps have been useful to have at least one contribution from a more cognitively orientated neuroscientist. There are many topics in neuropsychology that would have been sure to delight and inform the lay person about states of mind. Such topics might have included 'blindsight', spatial neglect, alien-hand syndrome, utilization behaviour, theory of mind, hallucinations, delusions and phantom limbs, all of which have been inspirational for a number of eminent neuroscientists and are probably susceptible to chemical modulation, though admittedly cannot yet be explained in terms of molecules.

The neuroendocrinological and neuro-immunological concomitants of stressful states are particularly well emphasized, although in such a concise book there is perhaps a little too much overlap (to the extent that both use a similar figure of the hypothalamic–pituitary axis) in two otherwise complementary chapters by McEwen and Sternberg. They also relate to a general theme of the biological bases of individual differences introduced in molecular-genetic (Hyman), developmental (Kagan) and psychiatric (Jamison) contexts. The latter two have a good balance of scientific, anecdotal and historical material. This is the type of book in which Galen and Byron feature more prominently in the index than Hubel or Bloom, but this is an eminently reasonable stratagem.

Overall, the style of the book is certainly seductive and the reader's hand is gently held by the editor during the introductory